

## INTERACTION OF ROOT-KNOT NEMATODE (*MELOIDOGYNE JAVANICA*) AND TOMATO AS AFFECTED BY HYDROGEN PEROXIDE

*Muwaffaq Ramadan Karajeh\**

Department of Plant Protection and IPM, Faculty of Agriculture, Mutah University  
Karak P.O. Box 7 (zip code 61710), Jordan

*Received: November 29, 2007*

*Accepted: May 05, 2008*

**Abstract:** The effect of hydrogen peroxide ( $H_2O_2$ ) on root-knot nematode (RKN, *Meloidogyne javanica*) in tomato was investigated. Soil drenching with exogenous  $H_2O_2$  was done using seven  $H_2O_2$  concentrations (1, 10, 100, 250, 500, 750 and 1000 mM) at different application times (24 hours before the time of plant inoculation with the RKN (T1), at the time of inoculation (T2), and 24 hours after the inoculation time (T3). The nematode reproduction rate (eggs/g fresh root) was significantly reduced in all  $H_2O_2$  treatments compared with the untreated control. The lowest reduction in nematode reproduction occurred at 10 mM  $H_2O_2$ . The application times T1 and T2 were significantly higher in reducing the reproduction rate than T3 at 250, 750 and 1000 mM  $H_2O_2$ . The content of endogenous  $H_2O_2$  in the treated plants was significantly higher than in the non-treated plants. Some phytotoxicity was apparent at the higher concentrations of  $H_2O_2$  ( $\geq 500$  mM) in the treated plants due to the accumulation of the endogenous  $H_2O_2$ . The treatments with 1 and 10 mM  $H_2O_2$  did not differ from the untreated control in plant chlorophyll content while the content was significantly reduced at the higher concentrations. Exogenous application of  $H_2O_2$  may have a direct effect on the nematode reproduction and an indirect effect on the treated tomato plants that can be elicited by  $H_2O_2$  to resist the nematode infection.

**Key words:** Host resistance, nematode reproduction, host-pathogen interaction.

### INTRODUCTION

Root-knot nematodes (RKNs, *Meloidogyne* spp.) attack a wide range of crop species. Annually, about 5% of the world crop production is destroyed by *Meloidogyne* species (Sasser *et al.* 1983; Barker *et al.* 1985; Sasser 1987).

---

\*Corresponding address:  
muwaffaq@mutah.edu.jo

Three *Meloidogyne* species (*M. javanica* (Teurb) Chitwood, *M. incognita* (Kofoid and White) Chitwood, race 1 and 2, and *M. arenaria* (Neal) Chitwood, race 2) were reported to occur in Jordan, with a predominance of *M. javanica* (Abu-Gharbieh 1982; Atieh 1986; Karajeh *et al.* 2005). The average annual losses of irrigated vegetable crops cultivated in the Jordan Valley due to RKNs has been estimated for nearly 15% (Abu-Gharbieh 1994)

Hydrogen peroxide ( $H_2O_2$ ) functions as a stress signal in plants, mediating adaptive responses to various stresses. Exposure to various abiotic and biotic stresses results in the accumulation of  $H_2O_2$  (Desikan *et al.* 2001). It is already known that  $H_2O_2$  can induce the expression of genes involved in antioxidant defense (Levine *et al.* 1994; Karpinski *et al.* 1999; Morita *et al.* 1999; Lopez-Huertas *et al.* 2000; Jaiti *et al.* 2004). Many studies have reported the killing capacity of  $H_2O_2$  produced by some bacteria or applied exogenously on nematodes e.g. *Caenorhabditis elegans* (Gustin *et al.* 2002; Jansen *et al.* 2002; Bolm *et al.* 2004).

To our knowledge, no previous studies were conducted to explore the effect of  $H_2O_2$  on plant-parasitic nematodes except for its use for clearing cyst walls and for increasing egg hatching when it is applied with irrigation to increase soil aeration (Goodey 1963). Hydrogen peroxide could affect the nematodes directly by its toxicity and/or indirectly as an elicitor triggering the host-plant defense. Therefore, this study was done to investigate the influence of  $H_2O_2$  applied exogenously on *M. javanica* and its interaction with tomato.

## MATERIALS AND METHODS

### Source of RKN Inoculum

Inoculum of *M. javanica* was obtained from a population that was naturally infecting eggplant plants in the central part of Jordan Valley. Galled root samples were washed thoroughly under tap water for 5 minutes. Small-galled roots were excised and examined under a dissecting microscope for finding developmental stages of the RKNs. Identification of *M. javanica* was done by observing female perineal patterns and measuring length of second-stage juveniles (Barker *et al.* 1985) and confirmed by species-specific SCAR-PCR (Karajeh 2004). The nematode eggs were extracted from galled roots by using 0.5% sodium hypochlorite (Hussey and Barker 1973).

### Plant Material and Treatments

Seeds of tomato cultivar Speedy were sown in nursery polystyrene trays filled with a pasteurized mixture of peatmoss, perlite and clay soil (1:2:1) in the greenhouse. The tomato plants were transplanted to pots and were transferred to a growth chamber ( $25 \pm 3^\circ C$  air temperature and 16 h day). Twenty-nine treatments were arranged in a completely randomized design with 10 replicates. A technical grade of  $H_2O_2$  was used. The treatments were soil drenching with  $H_2O_2$  at 7 concentrations (1, 10, 100, 250, 500, 750 and 1000 mM) in plant rhizosphere at three different application times: 24 hours before the time of plant inoculation with the RKN (T1), at the time of inoculation (T2), and 24 hours after the inoculation (T3). Treatments were repeated on un-inoculated plants (NI) and some inoculated plants were not treated with  $H_2O_2$ , used as controls. The seedlings were inoculated with 3000 eggs of *M. javanica* per pot. The experiment was ended after sixty days. The nematode eggs were re-extracted

from treated plants by using 0.5% NaOCl. The number of eggs/g root fresh weight was then determined.

### Analysis of data

Data were statistically analyzed using the general linear model (GLM) procedure of the system of analytical statistics (SAS). Least significant difference (LSD) test was used for mean separation at 0.05 probability (Steel and Torrie 1986).

### Determination of H<sub>2</sub>O<sub>2</sub> Content

Plant content of endogenous hydrogen peroxide was determined five days after the exogenous application according to Velikova *et al.* (2000). Briefly, plant material (500 mg fresh weight) was homogenized in 2 ml trichloroacetic acid (TCA) solution (1 g/l). After centrifugation at 12000Xg for 15 minutes, 0.5 ml of the supernatant was added to the reaction mixture containing 0.5 ml of 10 mM potassium phosphate buffer (pH 7.0) and 1 ml of 1 mM KI. Absorbance was determined at 390 nm using a UV spectrophotometer (Helios Alfa, Thermo Electron Corporation, USA) and the amount of H<sub>2</sub>O<sub>2</sub> was evaluated using a standard curve under the same conditions.

### Determination of Chlorophyll Content

Chlorophyll content was determined at the end of the experiment according to Seřtak *et al.* (1971). Samples (30 mg of fresh leaves) were immersed in 5 ml of 96% ethanol at 80°C for 10 min to extract the pigments. The absorbance of extracts was measured at 663 and 647 nm using the UV spectrophotometer. The amount of chlorophyll (chlorophyll a + chlorophyll b) was evaluated using a standard curve determined under the same conditions.

## RESULTS AND DISCUSSION

Compared with the untreated control, the RKN reproduction, expressed by the number of eggs/g fresh root was significantly reduced in all H<sub>2</sub>O<sub>2</sub> treatments. The lowest reduction in RKN reproduction occurred at 10 mM H<sub>2</sub>O<sub>2</sub> concentration used (Fig. 1). Regarding the application time of exogenous H<sub>2</sub>O<sub>2</sub>, T1 and T2 were definitely significantly higher in reducing the nematode reproduction than T3 at 250, 750 and 1000 mM H<sub>2</sub>O<sub>2</sub>. The reduction in the RKN reproduction rate could be correlated with the direct killing effect of H<sub>2</sub>O<sub>2</sub> of the nematode eggs and hatched juveniles. Hydrogen peroxide vapor was reported to have the ability to kill the eggs of *C. elegans in vitro* (Gustin *et al.* 2002). Streptococcus bacteria are able to kill *C. elegans* and this killing is only mediated by hydrogen peroxide. Streptococci can produce sufficient amounts of hydrogen peroxide to kill *C. elegans*, with killing kinetics similar to those of equimolar concentrations of pure hydrogen peroxide (Jansen *et al.* 2002; Bolm *et al.* 2004).

Content of endogenous H<sub>2</sub>O<sub>2</sub> in plants increased at higher concentrations of exogenous H<sub>2</sub>O<sub>2</sub> (≥ 500 mM) in treated plants compared with the control plants regardless the time of application (Fig. 2). There were significant differences between the treatments with exogenous H<sub>2</sub>O<sub>2</sub> used and the untreated control of uninoculated plants in the content of endogenous H<sub>2</sub>O<sub>2</sub> (Fig. 2). In general, chlorophyll content of treated plants was reduced with the increase in the concentration of exogenous H<sub>2</sub>O<sub>2</sub> used except for 1 and 10 mM where there was no difference (Fig. 3). At the higher concen-

tration of exogenous  $H_2O_2$ , treated plants were yellowish (low in chlorophyll content); some plants were stunted with relatively small root systems and some plants died before the end of the experiment. No significant differences were found among T1, T2 and T3 in plant contents of endogenous  $H_2O_2$  and chlorophyll (Fig. 2 and 3).

The use of exogenous  $H_2O_2$  at low concentrations (1–100 mM) did not affect plant growth. As a result of the application of exogenous  $H_2O_2$ , there were elevated levels of endogenous  $H_2O_2$  in the treated plants. Hydrogen peroxide is known as mobile and reactive compound in plants and in fact, tomato plants are known to accumulate relatively high levels of  $H_2O_2$  (Orozco-Cárdenas and Ryan 1999). Hydrogen peroxide has




Fig. 1. Effect of soil drenching with hydrogen peroxide on the reproduction of *Meloidogyne javanica* at three application times (DBI (T1): one day (24 hours) before the time of plant inoculation with the RKN, AI (T2): at the time of inoculation, and DAI (T3): one day (24 hours) after the inoculation). Columns followed by the same letters are not significantly different according to least significant (LSD) test at 0.05 probability level.




Fig. 2. Plant content of hydrogen peroxide five days after treatment with exogenous hydrogen peroxide at different concentrations in plants uninoculated (NI) or inoculated with *Meloidogyne javanica* (DBI (T1): one day (24 hours) before the time of plant inoculation with the RKN, AI (T2): at the time of inoculation, and DAI (T3): one day (24 hours) after the inoculation).  $H_2O_2$  content means ( $LSD_{0.05} = 0.25$ ) are not significant at 1, 10, 100 and 250 mM  $H_2O_2$  concentrations.




Fig. 3. Chlorophyll content of hydrogen peroxide five days after treatment with exogenous hydrogen peroxide at different concentrations in plants uninoculated (NI) or inoculated with *Meloidogyne javanica* (DBI (T1): one day (24 hours) before the time of plant inoculation with the RKN, AI (T2): at the time of inoculation, and DAI (T3): one day (24 hours) after the inoculation). Chlorophyll content means ( $LSD_{0.05} = 0.65$ ) are not significant at 1 and 10 mM  $H_2O_2$  concentrations.

been reported to play an important role in the development of disease resistance in plant species (Grant *et al.* 2000). Its accumulation in specific tissues, and in the appropriate quantities, is of benefit to plants and can mediate cross-tolerance toward other stresses (Bolwell 1999). The exogenous application of  $H_2O_2$  and ethephon protected tomato seedlings from chilling injury (Al-Haddad *et al.* 2002). The response to  $H_2O_2$  in tomato plant may be associated with the defense of plants against both herbivores and pathogens (Orozco-Cardenas and Ryan 1999). Hydrogen peroxide in plant cells has been shown to regulate the hypersensitive response and cell death as a defense response against pathogen attack (Klessig *et al.* 2000). At the higher concentrations of  $H_2O_2$  ( $\geq 500$  mM), the apparent toxicity in the treated plants might be due to the direct exposure of plant roots to the exogenous  $H_2O_2$  within few days after the application or due to the extreme accumulation of the endogenous  $H_2O_2$  in plant leaf tissues.

As a conclusion, the reduction of RKN reproduction may be attributed to the direct effect of the exogenous  $H_2O_2$  on the nematode eggs and hatched juveniles and indirectly to the  $H_2O_2$  effect on the treated tomato plants that could be elicited by  $H_2O_2$  to resist the nematode infection.

## ACKNOWLEDGEMENTS

This work was financially supported by the Scientific Research Deanship, Mutah University, Jordan.

## REFERENCES

- Abu-Gharbieh W.I. 1982. Distribution of *Meloidogyne javanica* and *M. incognita* in Jordan. *Nematol.* 28: 34–37.
- Abu-Gharbieh W.I. 1994. Root-Knot Nematodes, *Meloidogyne* spp. in Jordan: Biology and Control. Publication of University of Jordan, 2nd ed., Amman, Jordan, 97 pp.
- Al-Haddad J.M., Al-Jamali A.F. 2002. Effect of  $H_2O_2$  and ethephon spray on seedling chilling tolerance in three tomato cultivars. Poster No 667: Temperature stress. Annual meeting of the American Society of Plant Biologists, 3–7 August 2002, Denver, Colorado, USA.

- Atieh S. 1986. Pathogenicity and Histopathology of *Meloidogyne javanica* and *M. incognita* on Olive and Tomato. M. Sc. Thesis. Faculty of Agriculture, University of Jordan, 145 pp.
- Barker K.R., Sasser J.N., Carter C.C. 1985. An Advanced Treatise on *Meloidogyne*. Vol. II: Methodology. North Carolina State Univ. Graphics, Raleigh, NC, USA, 223 pp.
- Bolm M., Wouter T., Jansen M., Schnabel R., Chatwal G.S. 2004. Hydrogen peroxide-mediated killing of *Caenorhabditis elegans*: a common feature of different streptococcal species. *Infect. Immun.* 72: 1192–1194.
- Bolwell G.P. 1999. Role of active oxygen species and NO in plant defense responses. *Curr. Opin. Plant Biol.* 2: 287–294.
- Desikan R., Soheila A., Mackerness H., Hancock J.T., Neill S.J. 2001. Regulation of the *Arabidopsis* transcriptome by oxidative stress. *Plant Physiol.* 127: 159–172.
- Goodey J.B. 1963. Laboratory Methods for Work with Plant and Soil Nematodes. Ministry of Agriculture, Fisheries and Food, London, UK, Technical Bulletin. No. 2, 72 pp.
- Grant J.J., Yun B.W., Loake G.J. 2000. Oxidative burst and cognate redox signaling reported by luciferase imaging: identification of a signal network that functions independently of ethylene, SA and Me-JA but is dependent on MAPKK activity. *Plant J.* 24: 569–582.
- Gustin E.J., McDonnell G.E., Mullen G., Gordon B.E. 2002. The efficacy of vapor phase hydrogen peroxide against nematode infestation of the *Caenorhabditis elegans* model. 27–31 October 2002, 53rd AALAS National Meeting, San Antonio, Texas, USA.
- Hussey R.S., Barker K.N. 1973. A comparison of methods of collecting inocula of *Meloidogyne* spp. including a new technique. *Plant Dis. Report.* 57: 1025–1028.
- Jaiti F., Dihazi A., El-Hadrami A., El-Hassni M., El-Hadrami I. 2004. Effect of exogenous application of jasmonic acid on date palm defense reaction against *Fusarium oxysporum* f. sp. *albedinis*. *Phytopathologia Mediterranea* 43: 325–331.
- Jansen W.T., Bolm M., Balling R., Chhatwal G.S., Schnabel R. 2002. Hydrogen peroxide-mediated killing of *Caenorhabditis elegans* by *Streptococcus pyogenes*. *Infect. Immun.* 70: 5202–5207.
- Karajeh M.R. 2004. Identification, Distribution, and Genetic Variability of the Root-knot Nematodes (*Meloidogyne* spp.) in Jordan. Ph.D. Thesis, University of Jordan, Amman, Jordan, 152 pp.
- Karajeh M.R., Abu-Gharbieh W.I., Masoud S.H. 2005. First report of the root-knot nematode *Meloidogyne arenaria* Race 2 from several vegetable crops in Jordan. *Plant Dis.* 89, p. 206.
- Karpinski S., Reynolds H., Karpinska B., Wingsle G., Creissen G., Mullineaux P. 1999. Systemic signaling and acclimation in response to excess excitation energy in *Arabidopsis*. *Science* 284: 654–657.
- Klessig D.F., Durner J., Noad R., Navarre D.A., Wendehenne D., Kumar D., Zhou J.M., Shah J., Zhang S., Kachroo P. 2000. Nitric oxide and salicylic acid signaling in plant defense. *Proc. Natl. Acad. Sci.* 97: 8849–8855.
- Levine A., Tenhaken R., Dixon R., Lamb C. 1994. Hydrogen peroxide from the oxidative burst orchestrates the plant hypersensitive disease resistance response. *Cell* 79: 583–593.
- Lopez-Huertas E., Charlton W.L., Johnson B., Graham I.A., Baker A. 2000. Stress induces peroxisome biogenesis genes. *EMBO* 19: 6770–6777.
- Morita S., Kaminaka H., Masumura T., Tanaka K. 1999. Induction of rice cytosolic ascorbate peroxidase RNA by oxidative stress: the involvement of hydrogen peroxide in oxidative stress signaling. *Plant Cell Physiol.* 40: 417–422.
- Orozco-Cárdenas M.L., Ryan C.A. 1999. Hydrogen peroxide is generated systemically in plant leaves by wounding and systemin via the octadecanoid pathway. *Proc. Natl. Acad. Sci.* 96: 6553–6557.
- Sasser J. 1987. A perspective on nematode problems worldwide. p. 1–12. In: *Proceedings Nematodes Parasitic to Cereals and Legumes in Temperature Semiarid Regions*, Larnaca, Cyprus. ICARDA, Aleppo, Syria.

- Sasser J., Eisenback J., Carter C., Triantaphyllou A. 1983. The International *Meloidogyne* Project: Its Goals and Accomplishments. Annu. Rev. Phytopathol. 21: 271–288.
- Sešťák Z., Catsky J., Jarvis P. 1971. Plant Photosynthetic Production, Manual of Methods. Dr Junk Publishers, The Hague, Netherlands.
- Steel R.G., Torrie J.H. 1986. Biostatistics: Principles and Procedures. 3rd ed. McGraw-Hill Book Company, Inc., New York, USA, 662 pp.
- Velikova V., Yordanov I., Edereva A. 2000. Oxidative stress and some antioxidant systems in acid rain-treated bean plants. Protective role of exogenous polyamines. Plant Sci. 151: 59–66.

## POLISH SUMMARY

### WSPÓLDZIAŁANIE MATWIKA KORZENIOWEGO *MELOIDOGYNE JAVANICA* I ROŚLIN POMIDORA WYWOŁANE NADTLENKIEM WODORU

Badano wpływ nadtlenu wodoru ( $H_2O_2$ ) na porażenie korzeni pomidora mątwikiem *Meloidogyne javanica*. W tym celu nasączono ziemię roztworami nadtlenu wodoru o zróżnicowanym stężeniu (1, 10, 100, 250, 500, 750 i 1000 mM  $H_2O_2$ ) w trzech terminach: T1 – 24 h przed inokulacją patogenem; T2 – w czasie inokulacji; T3 – 24 h po inokulacji. We wszystkich traktowanych kombinacjach uzyskano istotne zmniejszenie reprodukcji mątwika (jaja/g świeżej masy korzeni) w porównaniu do nietraktowanej kombinacji kontrolnej. Najniższy stopień zmniejszenia reprodukcji nicienia stwierdzono przy użyciu stężenia 10 mM  $H_2O_2$ . Przy stosowaniu stężeń 250, 750 i 1000 mM  $H_2O_2$  w terminach T1 i T2 wystąpiło istotnie wyższe ograniczenie tempa reprodukcji szkodnika, niż w terminie T3. Zawartość endogennego  $H_2O_2$  w traktowanych roślinach była istotnie niższa niż w roślinach nietraktowanych. W przypadku wyższych stężeń ( $\geq 500$  mM  $H_2O_2$ ) stwierdzono na traktowanych roślinach fitotoksyczność, co było wynikiem nagromadzenia endogennego  $H_2O_2$ . W kombinacjach za stężeniami 1 i 10 mM  $H_2O_2$  rośliny nie różniły się pod względem zawartości chlorofilu od roślin w kombinacji kontrolnej, podczas gdy przy użyciu stężeń wyższych zawartość chlorofilu była istotnie niższa. Stwierdzono, że zewnętrzne zastosowanie  $H_2O_2$  może wywierać bezpośrednie działanie na reprodukcję *M. javanica*, oraz pośrednie działanie na rośliny pomidora, które stają się mniej wrażliwe na zakażenie tym patogenem.