

ORIGINAL ARTICLE

Do combinations of insecticides and acaricides influence spray droplet formation and the interaction with citrus leaves?

Jaqueline Franciosi Della Vechia*, Renata Thaysa Santos, Fabiano Griesang, Cícero Mariano Santos, Marcelo Costa Ferreira, Daniel Junior Andrade

Department of Crop Protection, São Paulo State University, São Paulo, Brazil

Vol. 59, No. 4: 487–493, 2019

DOI: 10.24425/jppr.2019.131257

Received: January 21, 2019

Accepted: May 15, 2019

*Corresponding address:

jaque_dellavechia@hotmail.com

Abstract

In agriculture, the mixing of pesticides in tanks is a common practice. However, it is necessary to pre-empt possible physical-chemical implications of this practice, which may affect the efficiency of the treatments performed. Therefore, the objective of this study was to evaluate the effects of the addition of acaricide to insecticidal spray mixtures on the formation of spray droplets and the interaction with citrus leaves. The experimental design was totally randomized, in a $(2 \times 3 + 1)$ factorial scheme for seven treatments. Factor A corresponded to the spray mixture used (isolate or in the mixture). Factor B corresponded to the insecticides tested (lambda-cyhalothrin + thiamethoxam, phosmet, and imidacloprid) and the control consisted of a spray mixture with spirotetramat only. Nine replications were performed for characterization of the spray droplet size spectrum and four replications for the analysis of the surface tension and the contact angle. The mixture of pesticides showed positive results in terms of application safety. The addition of acaricide to insecticide spray mixtures reduced the surface tension and contact angle of droplets on the adaxial surface of orange leaves. There was an increment in volume median diameter (VMD), a significant reduction in the volume of droplets with drift-sensitive size and improvement in the uniformity of droplet size. Therefore, the addition of acaricide to an insecticide spray mixture positively influenced spray droplet formation and the interaction with citrus leaves providing better coverage and droplet size fractions with an appropriate size for safe and efficient application.

Keywords: contact angle, coverage efficiency, droplet size, surface tension, tank mixture

Introduction

Citrus crops face phytosanitary problems related to the occurrence of disease vectors (insects and mites), which lead to reduced productivity and longevity of orchards, and depreciation of fruit quality. Among vector-borne diseases, citrus leprosis and HLB-Huanglongbing, transmitted by the mites *Brevipalpus* spp. and by the psyllid *Diaphorina citri* Kuwayama, 1908 (Bastianel *et al.* 2010; Bassanezi *et al.* 2013) are responsible for the greater demand for phytosanitary treatments and the consequently higher citrus production costs (Neves *et al.* 2002; Belasque Jr. *et al.* 2010; Van Leeuwen *et al.* 2015).

The application of pesticides by spraying is the main strategy adopted by farmers to keep the population of these vectors at low densities and the mixture of products is a common practice since these pests can occur at the same time in the field (Parra *et al.* 2010; Laranjeira *et al.* 2015). However, the combination of pesticides in a tank-mix can modify the physicochemical characteristics of the spray mixture and affect the values of the surface tension, pH, electrical conductivity and spray mixture stability. These parameters can, in turn, influence the efficacy of phytosanitary treatments (Maciel *et al.* 2010; Petter *et al.* 2013).

The physicochemical properties of the spray mixture can significantly change the process of droplet formation by changing their diameter and uniformity in size (Marmottant and Villermaux 2004), which influences the destination of droplet deposit. The droplet size is significant regarding susceptibility to drift and evaporation, penetration capacity, in addition to the deposit, spreading of the droplets, and distribution on the targets (Prokop and Kejklicek 2002; Andrade *et al.* 2010).

Studies on the interference of the physicochemical properties of the tank-mix on the formation, transport, and deposit of droplets on targets are essential because these investigations can allow for the evaluation of the efficiency and safety of the application (Hewitt 2008). Concerns about the interactions between products in sprayer tanks have been increasing among farmers, but there is a lack of information about associated consequences. Therefore, there is a gap between the stages of development and appropriate usage.

The objective of this work was to evaluate the effects of the addition of acaricide to insecticidal spray mixtures on the surface tension, droplet spectrum characteristics, as well as the contact angle of the droplets on the surface of citrus leaves.

Materials and Methods

The work consisted of three complementary experiments based on the analysis of surface tension, the contact angle of droplets on the surface of orange leaves [*Citrus sinensis* (L.) Osbeck] and the characterization of the spray droplet spectrum.

The spray liquids consisted of lambda-cyhalothrin + thiamethoxam [15 ml · 100 l⁻¹ (Engeo™ Pleno SC – Syngenta, India and Huddersfield, England)], phosmet [150 g · 100 l⁻¹ (Imidan® 500 WP – Cross Link, Brazil)], imidacloprid [20 ml · 100 l⁻¹ (Provado® 200 SC – Bayer CropScience AG, Germany)] and the acaricide spiroticlofen [25 ml · 100 l⁻¹ (Envidor® 240 SC; Bayer CropScience AG, Germany)], as a control. The spray mixtures were prepared with deionized water at an ambient temperature of 25 ± 2°C and a relative humidity of 65 ± 5%.

The experiments were performed in a completely randomized design in a 2 × 3 + 1 factorial scheme, with nine repetitions for the characterization of the spray droplet spectrum and four repetitions for the analyses of surface tension and contact angle. Factor A corresponded to the spray mixture used (insecticides isolated or mixed with acaricide). Factor B corresponded to the insecticides tested (lambda-cyhalothrin + thiamethoxam, phosmet, and imidacloprid) and the control consisted of a spray mixture with only spiroticlofen.

The mix of products was performed in the following order of preparation: water addition + solid formulation addition + liquid formulation addition, following NBR 13875 published in ABNT (2015).

Surface tension and droplet contact angle

Measurements of both parameters were performed in the Nucleus of Study and Development at the Technology of Application laboratory with an automatic tensiometer (OCA-15 Plus, Dataphysics Germany), following the methodologies of Decaro JR *et al.* (2015) and Ferreira *et al.* (2013). For surface tension measurements, the droplets were formed at the end of a needle coupled to a precision syringe of 500 µl, which was evaluated by the pendant drop method. In this approach, a CCD (charge-coupled device) high speed, high definition camera was used to capture images, and software was used to analyze the drop image by axis asymmetry (ADSA – Axisymmetric Drop Shape Analysis). The surface tension was determined using the Young-Laplace equation. The 20 sec data generated after drop formation was considered to characterize the tested spray mixture with respect to the inflection point of the dynamic surface tension curve. The droplet volume used during the surface tension measurement was 4 µl.

To determine the contact angle, droplets were deposited on the surface of orange leaves. In the process of acquiring droplet contact angles, longitudinal 5 × 1 (cm) rectangles of orange leaves were cut and fixed in a press so that any undulations on the analyzed surface would not interfere with the readings. The droplet volume used during the measurements of the contact angle was 2 µl. The evaluation of surface tension and droplet contact angle was carried out in a room with a controlled ambient temperature of 25 ± 2°C and relative humidity of 65 ± 5%.

Characterization of the spray droplet spectrum

To perform spraying, a hollow cone spray nozzle was used (TXA 8002, TeeJet®). Droplet size spectrum analysis was performed directly based on a laser beam diffraction method using a particle sizer (Mastersizer S®, Malvern Instruments Co.), well described by Griesang *et al.* (2017). It is based on the diffraction of light by the droplets, wherein the droplet diameter is inversely proportional to the angle of deviation of the light (Fernandes *et al.* 2007).

The spray nozzle was installed 40 cm away from the laser beam and the spray was driven by compressed air at a pressure of 0.5 MPa, which was held constant with a precision air pressure regulator. The spray liquids were prepared with deionized water in an

ambient temperature of $25 \pm 2^\circ\text{C}$ and relative humidity of $65 \pm 5\%$. The ambient conditions during the experiments were: an air temperature of 26°C , a relative humidity of 55% and the absence of wind.

To characterize the droplet size spectrum, the values of the volume median diameter (*VMD*), the percentage of the volume with droplet diameters less than $100 \mu\text{m}$ ($V < 100$) and the uniformity of the droplets (*SPAN*) were used. The *SPAN* was obtained from the droplet sizes that related to the spray volume at 10, 50 and 90% ($Dv0.1$, $Dv0.5$, and $Dv0.9$), using the following equation (Malvern 1997):

$$SPAN = \frac{Dv0.9 - Dv0.1}{Dv0.5}$$

The *SPAN* value indicates the variation of the droplet size i.e., the smaller the value, the more uniform the spray droplet spectrum.

Statistical analysis

The data for the surface tension, contact angle, *VMD*, $V < 100$, and the *SPAN* were subjected to analysis of variance by the *F* test and the means were compared using the Tukey test ($p > 0.05$), with the statistical program AgroEstat.

Pearson's correlation test was performed between the variables of surface tension and contact angle, *VMD* and between $V < 100$ and *SPAN*, at 5% probability.

Results

Surface tension

The addition of acaricide to an insecticide spray mixture reduced the surface tension ($F = 88.79$; $p < 0.0001$). There was a statistically significant difference between the insecticides when they were evaluated separately ($F = 4.75$; $p = 0.0198$). However, when combined with acaricide, the lambda-cyhalothrin + thiamethoxam insecticide showed a lower surface

tension value compared to the others. There was significant interaction between the spray mixture and insecticide factors ($F = 7.13$; $p = 0.0043$). Therefore, the factors can be considered interdependent (Table 1). The additional treatment (spirodiclofen) differed from the other treatments evaluated ($F = 366.14$; $p < 0.0001$), presenting lower values.

Droplet contact angle

Differences were observed among the analyzed factors regarding the contact angle of droplets on the adaxial surface of the orange leaves (Table 2). The addition of acaricide to the insecticidal spray mixtures reduced the contact angle of the droplets. Among the insecticides, imidacloprid showed a greater contact angle compared to the others. The contact angle of the control did not differ from other treatments ($F = 2.84$; $p = 0.1068$), with a value of 84.09.

The correlation coefficient was significant ($p = 0.0279$) and positive for the surface tension and contact angle for isolated insecticides ($R = 0.6307$). However, it was not significant ($p = 0.6565$) for the mixture of insecticide and acaricide ($R = 0.1434$). The addition of acaricide to insecticides reduced the dependence between factors.

Characterization of the spray droplet spectrum

Both the *VMD*, the $V < 100$, and the *SPAN* were influenced by the spray mixture constitution. The *VMD* values of the droplets presented significant differences for isolated insecticide spray mixtures or in a mixture with spirodiclofen ($F = 53.24$; $p < 0.0001$), and for the insecticides used ($F = 16.02$; $p < 0.0001$). The combination of insecticide and acaricide resulted in increases in the *VMD* value, except for insecticide imidacloprid (Fig. 1). For the insect-only spray mixture, imidacloprid presented the highest value of *VMD*. Among the insecticides combined with acaricide, phosmet presented a higher *VMD* value ($134.57 \mu\text{m}$). It was observed that the interaction between the evaluated factors was

Table 1. Mean values of surface tension coefficients of tested drops in $\text{mN} \cdot \text{m}^{-1}$ of spray mixture containing insecticide alone or in mixture with acaricide, related to the unfolding of degrees of freedom between the interactions of spraying liquid with insecticides

Insecticides	Spraying liquid	
	alone	mixture
Lambda-cyhalothrin + thiamethoxam	72.26 aA	67.42 bB
Phosmet	71.90 aA	70.33 bA
Imidacloprid	72.91 aA	68.96 bAB

LSD line (5%) 1.32; LSD column (5%) 1.60; CV (%) 1.29

Means followed by different lowercase letters in the lines and upper case in the columns differ significantly from each other by the Tukey's test ($p > 0.05$). LSD – least significant difference, CV – coefficient of variation

Table 2. Analysis of variance and tests of significance for the contact angle (degree) formed by drops applied on the adaxial surface of orange leaves in a function of different factors (spray mixture and insecticides)

Causes of variation	<i>F</i>	<i>p</i>
Spraying liquid (A)	5.71*	0.0263
Insecticides (B)	24.44**	< 0.0001
A vs. B	0.38 ns	0.6856
Control vs. Factorial	0.34 ns	0.5664
CV	4.73	

Spray mixture	
Alone	84.77 a
Mixture	80.94 b
LSD (5%)	3.33

Insecticides	
Lambda-cyhalothrin + thiamethoxam	78.74 b
Phosmet	79.05 b
Imidacloprid	90.78 a
LSD (5%)	4.95

Means followed by * or ** differ significantly from each other by the Tukey test, $p > 0.01$ or $p > 0.05$, respectively. Means followed by different lowercase letters in the columns differ significantly from each other by the Tukey test ($p > 0.05$). LSD – least significant difference; CV – coefficient of variation; ns – not significant




Fig. 1. (A) – Volumetric median diameter (*VMD*), (B) – percentage of the volume of droplets smaller than 100 μm ($V < 100$) and (C) – coefficient of droplet size uniformity (*SPAN*) for insecticides lambda-cyhalothrin + thiamethoxam (LT), phosmet (PM) and imidacloprid (IM) isolated and combined with acaricide applied with a hollow cone nozzle (TXA 8002). Columns followed by lowercase letters were used to compare insecticide alone and mixed with acaricide, while uppercase letters were used to compare between insecticides for alone and mixed spray separately. The same letters do not differ significantly from each other by the Tukey's test ($p > 0.05$)

significant ($F = 14.83$; $p < 0.0001$). The additional treatment did not differ from the other treatments, with the value 126.83 μm ($F = 1.31$; $p = 0.2567$).

The $V < 100$ differed between the isolated and mixed products ($F = 61.90$, $p < 0.0001$) and between the different insecticides ($F = 22.55$; $p < 0.0001$). There were significant differences in the interaction between the evaluated factors ($F = 7.27$; $p = 0.0016$) (Fig. 1). The spray mixture formed by the insecticide only presented a higher percentage of drops smaller than 100 μm in relation to the spray mixture combined with acaricide. Regardless of the type of spray mixture for the different insecticides, lambda-cyhalothrin + thiamethoxam presented a higher $V < 100$. The additional treatment (acaricide control) did not differ from the

other treatments ($F = 3.66$, $p = 0.0609$), with 26.42% of $V < 100$.

The uniformity of the droplet size spectrum was altered by the presence of acaricide ($F = 53.51$; $p < 0.0001$) and by the different insecticides evaluated ($F = 41.13$; $p < 0.0001$) and the results resembled those obtained for the variable $V < 100$. There was significant interaction between factors A and B ($F = 5.68$; $p = 0.0057$). Among the insecticides evaluated, the droplet size spectrum generated by lambda-cyhalothrin + thiamethoxam presented the least uniform droplet spectrum regardless of the type of spray mixture. The uniformity of the additional control (acaricide) differed from the other treatments ($F = 6.11$; $p = 0.0165$), presenting a value of 0.95.

The correlation coefficient was significant and negative between *VMD* and $V < 100$ for both isolated insecticides ($p = 0.0143$; $R = -0.6831$) and the insecticide and acaricide mixture ($p = 0.011$; $R = -0.7012$).

The Pearson correlation coefficient was significant ($p = 0.0206$) and positive for the isolated insecticides ($R = 0.6559$) between the parameters $V < 100$ and *SPAN*, and there was no significant correlation between the insecticide and acaricide ($p = 0.0947$; $R = 0.5041$).

Discussion

Surface tension and droplet contact angle

The mixing of pesticides in tanks intended for different uses is a current practice which is mainly used to save operational resources. However, it is necessary to know the possible physical, chemical and biological implications of this practice that could compromise the efficacy of the treatments performed (Andrade *et al.* 2013).

In general, we observed that the spray mixture surface tension was reduced with the addition of the acaricide, from 72.91 to 67.42. This is an indication of the spreading of droplets on the tested leaves and the greater coverage of the leaf's surface. A positive correlation was recorded as has been observed in other research (Yu *et al.* 2009; Decaro JR *et al.* 2014). This effect may be related to the modification of the molecular orientation in the active ingredients, or the interaction of the liquid and the leaf's surface (Iost and Raetano 2010).

For insecticide spraying of only the tested mixtures, the values were similar to that obtained for water ($74 \text{ mN} \cdot \text{m}^{-1}$). This implies that this kind of spray mixture has high surface tension values and the droplets tend to maintain high contact angles on the surface where they settle. When the contact angle exceeds 90° , the surface is considered as hydrophobic, whereas lower values characterize the surface as hydrophilic (Iost and Raetano 2010). The constituents of the leaf's epidermis have a significant influence on this spreading, and when the interfacial tension (spray mixture-leaf) is greater than the surface tension of the liquid, it results in greater spreading (Damak *et al.* 2016).

The coverage provided by the tested spray mixtures is supported by the values of the surface tension since the Pearson correlation coefficient between these factors was positive. The cuticle of the orange leaves contains wax, which is a hydrophobic substance, and the addition of products with tenso-active properties in the spray mixture can provide a greater affinity with this substance, resulting in greater spreading (Iost and Raetano 2010).

In a field application, high surface tensions and contact angles would imply a lower potential for coverage of the target. This would require an increase in the application volumes or the use of smaller droplets. Considering that orange crop plants usually require coverage of a large leaf surface, the volume requirement is already high. For targets of low mobility and difficult access to spraying, such as *Brevipalpus* spp. mites, the use of spray mixtures with high surface tensions and contact angles can, therefore, result in more expensive applications (Decaro JR *et al.* 2015).

Thereby, the adoption of alternatives that improve the coverage of the treated surface, such as the use of adjuvants in the spray mixtures as the products evaluated in this study, can contribute to the satisfactory coverage of the targets without the need to increase the volume of application. This characteristic can have direct implications on the costs of phytosanitary treatments due to the operational performance of the sprayer by using less water per area unit. However, the costs of the adjuvants should not be greater than the operational costs provided by the alternative of application with higher volumes. In addition, alternatives that modify the characteristics of the spray mixture may result in a reduction of the expected biological effect for the control of the targets.

Characterization of spray droplet size

The composition of phytosanitary spray mixtures based on multiple components can influence the viscosity and surface tension of the spray mixture by acting on the internal structure of the jet, with an overall effect on the size of the droplets formed (Butler-Ellis *et al.* 2001). The addition of acaricide to the insecticidal spray mixtures resulted in increases of *VMD* values, *SPAN* reduction and $\% \text{vol} < 100 \mu\text{m}$ with combinations, compared to isolated products. Generally, the increase in droplet size leads to less coverage of leaves in the field. However, with the increased spread of the droplets and the reduction in the volume of droplets susceptible to drift, the combination of products can produce positive results.

The uniformity of the droplet size spectrum determined with *SPAN* can be altered based on the composition of the spray mixture (Costa *et al.* 2017). In this way, the addition of acaricide to the insecticides can facilitate a more homogenous droplet size spectrum. According to the values obtained for the Pearson correlation coefficient, there was a negative correlation between the *VMD* and $V < 100$ factors for both the isolated insecticides and the mixture with the acaricide. Therefore, with an increase of the *VMD* values, there was a decrease in the $V < 100$ (Baesso *et al.* 2014; Costa *et al.* 2017). An increase of the *VMD* or a reduction of the amount in droplets smaller than $100 \mu\text{m}$ can result

in less drift risk (Dorr *et al.* 2013; Oliveira *et al.* 2015). This can, in turn, reduce the risk of contamination in adjacent areas.

A positive correlation was found for the $V < 100$ and SPAN factors, meaning that a more uniform droplet size spectrum leads to a reduction in $V < 100$. Considering that droplets are classified according to sizes into several classes (ASABE 2009), the very fine class can suffer significant spray losses due to drift and evaporation. Droplets with a size greater than those recommended for the application can be lost by a ricochet and superficial runoff. Obtaining a jet of spray mixture with droplets of uniform size spectrum means lower losses (Ferreira *et al.* 2013). It is worth mentioning that the use of the class of fine droplets is recommended when greater coverage and penetration is required in the canopy of plants, as well as those produced in this research, and that classes of medium or coarse droplet sizes are better for application under conditions of greater risk of drift (Cunha *et al.* 2007; De Oliveira and Antuniassi 2011; Almeida *et al.* 2014, 2016; Triloff *et al.* 2014). However, little is known about the drop diameter ratio that provides coverage of a leaf's surface that results in effective pest control. The ideal coverage for the response to biological phenomenon will also depend on the interactions of the molecules and the formulation of the product with the target (Almeida *et al.* 2016).

Conclusions

The combinations of tested products showed notably positive results on the safety of application. A significant reduction of the drift susceptible fraction of droplets and an increase in the VMD value were observed. In addition, the improvement in the uniformity of the droplet size spectrum is worth mentioning, because it resulted in a larger fraction of droplet uniformity with the appropriate size for the application. In addition, a decrease in surface tension improved spreading of the drops on the target surface, resulting in increased coverage values and consequently greater control of biological targets.

References

- ABNT [Associação brasileira de normas técnicas]. 2015. NBR 13875: Agrotóxico – Avaliação de compatibilidade físico-química. 2nd ed. Rio de Janeiro, Brazil, 15 pp. (in Portuguese)
- Almeida D.P., Timossi P.C., Lima S.F., Silva U.R., Reis É.F. 2014. Weather conditions and application volumes in the desiccation of *Urochloa ruziziensis* and spontaneous vegetation. *Revista Brasileira de Herbicidas* 13: 245–251.
- Almeida D.P., Agostini A.R., Yamauchi A.K.F., Decaro Junior S.T., Ferreira M.C. 2016. Application volumes and sizes of droplets for the application of diquat herbicide in the control of *Eichhornia crassipes*. *Planta Daninha* 34 (1): 171–179. DOI: <http://dx.doi.org/10.1590/S0100-83582016340100018>
- Andrade D.J., Ferreira M.C., Fenólio L.G. 2013. Compatibilidade entre acaricidas e fertilizantes foliares em função de diferentes águas no controle do ácaro da leprose dos citros *Brevipalpus phoenicis*. *Revista Brasileira de Fruticultura* 35 (1): 39–50. DOI: <http://dx.doi.org/10.1590/S0100-29452013000100006> (in Portuguese, with English summary)
- Andrade D.J., Ferreira M.C., Santos N.C. 2010. Efeito da adição de óleos ao acaricida cyhexatin sobre o ácaro *Brevipalpus phoenicis* e na retenção de calda por folhas de citros. *Revista Brasileira de Fruticultura* 32 (4): 1055–1063. DOI: <http://dx.doi.org/10.1590/S0100-29452010000400014> (in Portuguese, with English summary)
- ASABE [American Society of Agricultural and Biological Engineers]. 2009. Spray nozzle classification by droplet spectra. p. 1–3. In: “ASABE Standard”. St. Joseph
- Baesso M.M., Teixeira M.M., Ruas R.A.A., Baesso R.C.E. 2014. Tecnologia de aplicação de agrotóxicos. *Revista Ceres* 61: 780–785. DOI: <http://dx.doi.org/10.1590/0034-737x201461000003> (in Portuguese)
- Bassanezi R.B., Montesino L.H., Gimenes-Fernandes N., Yamamoto P.T., Gottwald T.R., Amorim L., Bergamin Filho A. 2013. Efficacy of area-wide inoculum reduction and vector control on temporal progress of Huanglongbing in young sweet orange plantings. *Plant Disease* 97 (6): 789–796. DOI: <http://dx.doi.org/10.1094/PDIS-03-12-0314-RE>
- Bastianel M., Novelli V.M., Kitajima E.W., Kubo K.S., Bassanezi R.B., Machado M.A., Freitas-Astúa J. 2010. Citrus leprosis: centennial of an unusual mite-virus pathosystem. *Plant Disease* 94: 284–292. DOI: <http://dx.doi.org/10.1094/PDIS-94-3-0284>
- Belasque Júnior J., Bassanezi R.B., Yamamoto P.T., Ayres A.J., Tachibana A., Violante A.R., Tank J.R.A., Di Giorgi F., Terzi F.E.A., Menezes G.M., Dragone J., Jank Júnior R.H., Bové J.M. 2010. Lessons from Huanglongbing management in São Paulo State, Brazil. *Journal of Plant Pathology* 92 (2): 285–302. DOI: <http://dx.doi.org/10.4454/jpp.v92i2.171>
- Brasil. Ministério da Agricultura, Pecuária e Abastecimento. 2002. Decreto 4074 que regulamenta a Lei 7802 de 11 de julho de 1989 que dispõe sobre agrotóxicos, seus componentes e afins e de outras providências. *Diário Oficial da União, Brasília, DF*. (in Portuguese)
- Butler-Ellis M.C., Tuck C.R., Miller P.C.H. 2001. How surface tension of surfactant solutions influences the characteristics of sprays produced by hydraulic nozzles used for pesticide application. *Colloids Surfaces A: Physicochemical and Engineering Aspects* 190 (3): 267–276. DOI: [http://dx.doi.org/10.1016/S0927-7757\(00\)00776-7](http://dx.doi.org/10.1016/S0927-7757(00)00776-7)
- Costa L.L., Carneiro A.L.C.G., Souza A.D.V., Almeida D.P., Ferreira M.C. 2017. Caracterização da aplicação com diferentes inseticidas e pontas de pulverização na cultura da soja. *Engenharia na Agricultura* 25 (2): 116–123. DOI: <http://dx.doi.org/10.13083/reveng.v25i2.732> (in Portuguese, with English summary)
- Cunha J.P.A.R., Teixeira M.M., Fernandes H.C. 2007. Avaliação do espectro de gotas de pontas de pulverização hidráulicas utilizando a técnica da difração do raio laser. *Engenharia Agrícola* 27: 10–15. DOI: <http://dx.doi.org/10.1590/S0100-69162007000200002> (in Portuguese, with English summary)
- Damak M., Mahmoudi S.R., Hyder M.N., Varanasi K.K. 2016. Enhancing droplet deposition through *in-situ* precipitation. *Nature Communications* 7 (12560): 1–9. DOI: 10.1038/ncomms12560
- Decaro JR S.T., Ferreira M.C., Lasmar O., Campos H.B.N. 2014. Relationship among variables of sprays applied at reduced volumes in a coffee plantation. *Aspects of Applied Biology* 122 (1): 415–422.
- Decaro JR S.T., Ferreira M.C., Lasmar O., Fouquet G. 2015. Reducing spray volume for the control of *Leucoptera coffeella* (Lepidoptera: Lyonetiidae) in coffee plants. *Coffee Science* 10 (4): 491–498.

- De Oliveira M.A.P., Antuniassi U.R. 2011. Eficácia do flutriafol e do flutriafol + tiofanato metílico aplicados com gotas finas ou médias no controle da ferrugem asiática da soja. *Revista Energia na Agricultura* 26 (1): 94–112. DOI: <http://dx.doi.org/10.17224/EnergAgric.2011v26n1p94-112> (in Portuguese, with English summary)
- Dorr G.J., Hewitt A.J., Adkins S.W., Hanan J., Zhang H., Noller B. 2013. A comparison of initial spray characteristics produced by agricultural nozzles. *Crop Protection* 53: 109–117. DOI: <http://dx.doi.org/10.1016/j.cropro.2013.06.017>
- Fernandes A.P., Parreira R.S., Ferreira M.C., Romani G.N. 2007. Caracterização do perfil de deposição e do diâmetro de gotas e otimização do espaçamento entre bicos na barra de pulverização. *Engenharia Agrícola* 27 (3): 728–733. DOI: <http://dx.doi.org/10.1590/S0100-69162007000400016> (in Portuguese, with English summary)
- Ferreira M.C., Lasmar O., Decaro Junior S.T., Neves S.S., Azevedo L.H. 2013. Quality of insecticide application on peanut (*Arachis hypogaea* L.) with and without adjuvants in the spraying liquid, under simulated rain. *Bioscience Journal* 29 (5): 1431–1440.
- Griesang F., Decaro R.A., dos Santos C.A.M., Santos E.S., Lima Roque N.H., da Costa Ferreira M. 2017. How much do adjuvant and nozzles models reduce the spraying drift? Drift in agricultural spraying. *American Journal of Plant Science* 8: 2785–2794. DOI: <https://doi.org/10.4236/ajps.2017.811188>
- Hewitt A.J. 2008. Spray optimization through application and liquid physical property variables-I. *The Environmentalist* 28 (1): 25–30. DOI: <https://doi.org/10.1007/s10669-007-9044-5>
- Iost C.A.R., Raetano C.G. 2010. Tensão superficial dinâmica e ângulo de contato de soluções aquosas com surfatantes em superfícies artificiais e naturais. *Engenharia Agrícola* 30 (4): 670–680. DOI: <http://dx.doi.org/10.1590/S0100-69162010000400011> (in Portuguese, with English summary)
- Laranjeira F.F., Silva S.X.B., Andrade E.C., Almeida D.O., Silva T.S.M., Soares A.C.F., Freitas-Astúa J. 2015. Infestation dynamics of *Brevipalpus phoenicis* (Geijskes) (Acari: Tenuipalpidae) in citrus orchards as affected by edaphic and climatic variables. *Experimental and Applied Acarology* 66 (4): 491–508. DOI: <http://dx.doi.org/10.1007/s10493-015-9921-4>
- Maciel C.D.G., Guerra N., Oliveira Neto A.M., Poletine J.P., Bastos S.L.W., Dias N.M.S. 2010. Tensão superficial estática de misturas em tanque de glyphosate + chlorimuronethyl isoladas ou associadas com adjuvantes. *Planta Daninha* 28 (3): 673–685. DOI: <http://dx.doi.org/10.1590/S0100-83582010000300025>
- Malvern 1997. *Getting Started – Manual Mastersizer* (No. 1). 1.3, United Kingdom, 183 pp.
- Marmottant P., Villermaux E. 2004. On spray formation. *Journal of Fluid Mechanics* 498 (498): 73–111. DOI: <http://dx.doi.org/10.1017/S0022112003006529>
- Neves E.M., Dayoub M., Dragone D.S. 2002. Demanda por fatores de produção na citricultura: Fertilizantes e defensivos agrícolas. *Laranja* 23 (1): 37–56. (in Portuguese, with English summary)
- Oliveira R.B., Antuniassi U.R., Gandolfo M.A. 2015. Spray adjuvant characteristics affecting agricultural spraying drift. *Journal of the Brazilian Association of Agricultural Engineering* 35: 109–116. DOI: <https://doi.org/10.1590/1809-4430-Eng.Agric.v35n1p109-116/2015>
- Parra J.R.P., Lopes J.R.S., Torres M.L.G., Nava D.E., Paiva P.E.B. 2010. Bioecologia do vetor *Diaphorina citri* e transmissão de bactérias associadas ao huanglongbing. *Citrus Research & Technology* 31 (1): 37–51. DOI: <https://doi.org/10.5935/2236-3122.20100004> (in Portuguese, with English summary)
- Petter F.A., Segate D., Almeida F.A., Alcântara Neto F., Pacheco L.P. 2013. Incompatibilidade física de misturas entre inseticidas e fungicidas. *Comunicata Scientiae* 4 (2): 129–138. (in Portuguese, with English summary)
- Prokop M., Kejklicek R. 2002. Effect of adjuvants on spray droplet size of water. *Engineering in Agriculture* 48: 144–148.
- Triloff P., Lind K., Matzer W., Knoll M., Kleisinger S. 2014. Adjusting and straightening the air distribution of sprayers for three-dimensional crops: The state of the art. p. 133–138. In: *Proceedings of the 5th European Workshop on Standardized Procedure for the Inspection of Sprayers*. Montpellier, France.
- Van Leeuwen T., Tirry L., Yamamoto A., Nauen R., Dermauw W. 2015. The economic importance of acaricides in the control of phytophagous mites and an update on recent acaricide mode of action research. *Pesticide Biochemistry Physiology* 121: 12–21. DOI: <https://doi.org/10.1016/j.pestbp.2014.12.009>
- Yu Y., Zhu H., Ozkan H.E., Derksen R.C., Krause C.R. 2009. Evaporation and deposition coverage area of droplets containing insecticides and spray additives on hydrophilic, hydrophobic, and crabapple leaf surfaces. *American Society of Agricultural and Biological Engineers* 52 (1): 39–49. DOI: <https://doi.org/10.13031/2013.25939>