

ORIGINAL ARTICLE

Combinations of *Tagetes filifolia* Lag. essential oil with chemical fungicides to control *Colletotrichum truncatum* and their effects on the biocontrol agent *Trichoderma harzianum*

Laura Carolina Gadban¹, Boris Xavier Camiletti^{2,3*}, Ezequiel Darío Bigatton²,
Silvia Graciela Distéfano¹, Enrique Iván Lucini²

¹ Estación Experimental Agropecuaria Marcos Juárez/Laboratorio de Fitopatología, Instituto Nacional de Tecnología Agropecuaria (INTA), Córdoba, Argentina

² Microbiología Agrícola, Facultad de Ciencias Agropecuarias-Universidad nacional de Córdoba (UNC), Córdoba, Argentina

³ Instituto Multidisciplinario de Biología Vegetal (IMBIV), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Córdoba, Argentina

Vol. 60, No. 1: 41–50, 2020

DOI: 10.24425/jppr.2020.132202

Received: June 17, 2019

Accepted: August 7, 2019

*Corresponding address:
bcamiletti@agro.unc.edu.ar

Abstract

Soybean [*Glycine max* (L.)], one of the most important crops in Argentina, is commonly infected by *Colletotrichum truncatum*, the causal agent of anthracnose. *Tagetes filifolia* essential oil (EO) is presented as a natural approach to minimize the dose of chemical fungicides applied to the crop. The fungus *Trichoderma harzianum* is used as a biocontrol agent because of its ability to produce secondary metabolites that destroy cell walls of phytopathogenic fungi. However, its performance can be affected when it is exposed to chemical fungicides. The objective of this work was to evaluate the antifungal activity of *T. filifolia* EO both individually and combined with chemical fungicides against *C. truncatum*, and its effect on *T. harzianum*. Fungi were isolated from soybean crops. The following pesticides were assessed: carbendazim (F1), difenoconazole (F2) and trifloxystrobin + cyproconazole (F3). The EO was obtained from native plants and its chemical composition was analyzed by gas chromatography–mass spectrometry (GC–MS). The minimum fungicide concentration (MFC) was determined for each compound. Fungicides were combined with the EO to look for combinations that allowed a reduction of pesticide doses. Among fungicides, F1 showed the strongest antifungal activity against *C. truncatum* (MFC = 0.25 $\mu\text{l} \cdot \text{l}^{-1}$) and *T. harzianum* (MFC = 1.5 $\mu\text{l} \cdot \text{l}^{-1}$). The sensitivity of both fungi to the EO was lower than to fungicides. The EO presented MFCs of 6,000 and 9,000 $\mu\text{l} \cdot \text{l}^{-1}$ against *C. truncatum* and *T. harzianum*. The EO and F1 affected the growth of *T. harzianum* at concentrations that controlled *C. truncatum* (31 and 10%). Eight combinations of fungicides and the EO allowed fungicide concentration reductions of up to 80%, although the growth of the biocontrol strain was also affected. The results demonstrated that *T. filifolia* EO can be used to control anthracnose and reduce doses of chemical fungicides applied to soybean crops. Its effect on *T. harzianum* should be considered in the design of integrated pest management strategies.

Keywords: anthracnose, integrated pest management, natural control, soybean

Introduction

Soybean [*Glycine max* (L.)] is widely cultivated in many countries due to its well-developed agronomic performance, characterized by a great protein yield per unit area (Vollmann 2016). In Argentina, 20.5 million

hectares of this commodity were sown during the 2016–2017 growing season. Soybean reached an overall production of 70.2 million tons, representing 31% of total exports of the country (Phélinas and

Choumert 2017). Soybean production is highly affected by late season diseases (LS), a combination of several pathogens that may cause yield and productivity losses, as well as seed quality changes (Carmona *et al.* 2015). Anthracnose, mainly caused by *Colletotrichum truncatum*, is one of the most prevalent LS in Argentinean soybean production (Ramos *et al.* 2013). Infections with this fungus are characterized by twisted and aborted pods (Ramos *et al.* 2013; Carmona *et al.* 2015; Dias *et al.* 2016).

The use of chemical fungicides is a common practice employed in the control of soybean diseases on a global scale (Carmona *et al.* 2011; Carmona *et al.* 2015; Dias *et al.* 2016). Several synthetic fungicides that belong to benzimidazoles, strobilurins and triazoles groups have been extensively used in the field to control *C. truncatum* (Gawade *et al.* 2009; Carmona *et al.* 2011; Torres-Calzada *et al.* 2015). However, it is increasingly frequent to find isolates that are less sensitive or even resistant to these fungicides (Fernandez-Ortuño *et al.* 2012; Torres-Calzada *et al.* 2015). Triazole resistance among *Aspergillus* and *Candida* species has been well documented (Goncalves *et al.* 2016). Furthermore, chemical residues from fungicides can exhibit low degradability, remaining in soil and water for years. These are some of the reasons why chemical treatment is recommended as part of an integrated fungal control program (Sosa-Gomez *et al.* 2003).

Essential oils (EOs) are complex mixtures of volatile components synthesized as secondary metabolites by aromatic plants in order to protect themselves against infectious diseases (Isman 2000). These natural extracts may have numerous components, although in general, a few major compounds determine their biological properties (Isman 2000; Bakkali *et al.* 2008). Essential oils are easily degraded by non-target microorganisms, and are considered safe for humans and animals (Isman 2000). Numerous EOs, as well as their individual components, have been characterized by their antifungal activities against several plant pathogens (Abbaszadeh *et al.* 2014; Camiletti *et al.* 2014; Camiletti *et al.* 2016). Thus, the application of EOs has become a very attractive approach for the control of several phytopathogenic fungi (Omidbeygi *et al.* 2007; Bluma *et al.* 2008; Abdel-Kader *et al.* 2011; Camiletti *et al.* 2014; Camiletti *et al.* 2016). Generally, the sensitivity of fungal species to EOs is lower than to chemical fungicides. Therefore, EOs are combined with synthetic compounds in order to improve their performance, as well as to reduce pesticide doses (Pyun and Shin 2006; Lim and Shin 2008; Camiletti *et al.* 2016). To reach this objective, the antifungal activity of EOs is frequently investigated in combination with chemical pesticides in search of synergistic interactions (Shin and Kang 2003; Pyun and Shin 2006; Pavela 2014). However, additive or antagonistic interactions may also appear when two

antimicrobial substances are combined (Kosman and Cohen 1996). In previous studies, *Tagetes filifolia* EO was successfully combined with a synthetic fungicide to control *Sclerotium cepivorum* in garlic crops (Camiletti *et al.* 2016).

Species of the fungal genera *Trichoderma*, characterized by antagonistic properties against phytopathogenic fungi, are commonly used to control plant diseases. The success of *Trichoderma* species is mainly due to its ability to colonize different substrates, its effectiveness for nutrient use and strong aggressiveness against phytopathogenic fungi (Benítez *et al.* 2004). The most common strains utilized as biocontrol agents belong to the species *Trichoderma harzianum* (Benítez *et al.* 2004; Sushir *et al.* 2015; Ganuza *et al.* 2017). This fungus destroys cell walls of phytopathogenic fungi by production of lytic enzymes and secondary metabolites (Grondona *et al.* 1997; Sharma *et al.* 2017). *Trichoderma harzianum*, an ubiquitous species present in crops, soil, and debris, may act as a biocontrol agent against ascomycetous, deuteromycetous and basidiomycetous fungi (Benítez *et al.* 2004).

Integrated pest management commonly involves biological, physical and chemical strategies (Abrol and Shankar 2016). Some EOs have been successfully combined with biocontrol agents and constitute an applicable method for controlling plant diseases (Arrebola *et al.* 2010; Abdel-Kader *et al.* 2011). On the other hand, some antifungal compounds inhibit the growth of biocontrol microorganisms and may affect its performance (Sarkar *et al.* 2010). The objective of this work was to evaluate the antifungal activity of *T. filifolia* EO, both individually and combined with chemical fungicides, against *C. truncatum* and their effects on the biocontrol agent *T. harzianum*.

Materials and Methods

Plant material and essential oil extraction

Tagetes filifolia plants were grown in 3-liter pots containing soil collected from the experimental station of the National Institute of Agricultural Technology (INTA) in Marcos Juárez (Córdoba, Argentina). Pots were placed in a greenhouse where the temperature ranged from 25 to 28°C. Developing plants were transferred to field plots at the grown stage of two detectable nodes. Plants were watered every 7 days and harvested when 50% of the plants had at least one corolla open. Aerial parts were dried in a forced air oven at 60°C for 4 days and hydrodistilled for 2 h in a Clevenger-type apparatus coupled to a separated extraction chamber (Asensio *et al.* 2011). The essential oil was kept in dark flasks at -20°C and the yield was determined by measuring the collected volume per plant weight.

Essential oil chemical composition

The essential oil composition was determined by CG-MS in a Perkin Elmer Clarus 600 chromatograph coupled with an ion trap mass detector. Compounds were separated using a capillary column DB-5 (30 m, 0.25 mm internal diameter, and 0.25- μ m coating thickness) and helium as the carrier gas with a flow rate of 0.9 ml \cdot min⁻¹. Ionization was performed by electron impact at 70 eV and the mass spectral data were obtained in the scan mode in the m/z range 35 to 450. The temperature program was settled at 60°C for 5 min and from 60 to 250 at a rate of 5°C/min (Asensio et al. 2011). The compounds were identified by contrasting their retention time and mass spectra with published data (Adams 1995) and NIST libraries (version 3.0). Quantitative analysis was performed using the area normalization method and the results were expressed as relative percentages (Asensio et al. 2011).

Fungal isolates

Seeds, pods and stems that looked infected by *C. truncatum* were collected from a soybean crop cultivated in the Marcos Juarez area at the end of the growing season, immediately before harvest. Samples were surface-disinfected by dipping in ethanol solution (70%) and then washed three times in sterile water. Plant material was placed on Petri plates (9 cm) previously prepared with sterile moistened paper. Samples were placed in a culture chamber at 25 \pm 1°C for 4 days, alternating 12 h of light with 12 h of darkness. *Colletotrichum* acervuli were separated and examined under a stereo microscope. Spores of *C. truncatum* were identified according to morphological characteristics (Armstrong-Cho and Banniza 2006), transferred to Petri plates containing potato dextrose agar (Britania, Buenos Aires, Argentina) and incubated as previously described. To confirm identity, Petri plates were prepared with sterile soybean stems and inoculated with mycelium from a 7-day culture (Dhingra and Sinclair 1995). Samples were again incubated as described above. After incubation, acervuli were separated and the morphological characteristics of the spores were compared with those indicated for *C. truncatum* (Manandhar and

Hartman 1999; Armstrong-Cho and Banniza 2006). *Trichoderma harzianum* was isolated from soil samples collected from the same soybean crop. Soil samples (10 g) were suspended in physiological saline solution. Serial dilutions were prepared and seeded on Petri plates containing selective medium for *Trichoderma* (Dhingra and Sinclair 1995). Samples were incubated under the same conditions as *C. truncatum*. Isolates characterized as *T. harzianum* were sub-cultured on potato dextrose agar and incubated as previously described. The identity of the isolates was confirmed following the taxonomic schemes described by Rifai (1969). The distinctive features of *T. harzianum* are conidiophores which are compactly branched in a pyramidal shape, the base being highly branched, and the apex usually bearing a solitary phialide. Conidia are less than 3.5 μ m long, nearly spherical and smooth-walled (Rifai 1969).

Chemical fungicides

Fungicides included in this study were selected based on their current use to control anthracnose caused by *C. truncatum* in Argentinean soybean crops. Commercial formulations were purchased from local suppliers (Table 1). These products were used instead of active ingredients in an attempt to obtain a more accurate predictive value during the in vitro assay. The response of the pathogens was based on the formulation employed in the field.

Antifungal activity

A broth method was used to determine the antifungal activity of the EO and fungicides against *C. truncatum* and *T. harzianum* according to previous bibliography (Lucini et al. 2006; Camiletti et al. 2014). Petri plates were prepared with 20 ml of liquid growing medium (20% v/v lixiviated potato; 2% w/v glucose; pH 4.5). The EO and fungicide solutions were added to the liquid medium according to the desired concentration. Prior to addition, the EO and fungicides were diluted in ethanol and sterile water, respectively. The final concentration of ethanol was always less than 1% v/v. A 5-mm diameter agar disc was taken from 7-day-old fungal colonies and used to inoculate selected

Table 1. Fungicides included in the assays

Code	Group	Active ingredient [% w/v]	Formulation	Class*
F1	benzimidazoles	carbendazim (50)	SC	IV
F2	triazoles	difenoconazole (25)	EC	III
F3	strobilurins + triazoles	trifloxystrobin (32) + cyproconazole (14)	SC	IV

EC – emulsifiable concentrate, SC – suspension concentrate

*toxicological categories based on Argentinean standard regulations (CASAFE 2015): IV – product that normally does not represent danger, III – low dangerous product

treatments. Petri plates without the addition of an antifungal agent were included as a control. All treatments were incubated in a culture chamber for 7 days at $25 \pm 1^\circ\text{C}$ alternating 12 h of light with 12 h of darkness. After incubation, the diameter of the mycelium was measured to calculate the percentage of growth inhibition (*PGI*) using the following equation (Camiletti *et al.* 2014):

$$PGI = [(C - T)/C] \times 100,$$

where: *C* and *T* are the mean values ($n = 3$) of hyphal extension (mm) obtained for the control and treatments with antifungal agents, respectively. The lowest concentration of active ingredient or EO in which the *PGIs* were equal to 100% represented the minimum fungicidal concentration (*MFC*) values (Abbaszadeh *et al.* 2014; Camiletti *et al.* 2014).

Mixture effects

The broth method was also utilized to determine the interaction of different concentrations of fungicides with the EO. The selected treatments were composed of all possible combinations between 20, 40, 60 or 80% of the previously determined *MFC* of EO and 20, 40, 60 or 80% of the previously determined *MFCs* of fungicides (Pyun and Shin 2006; Camiletti *et al.* 2016). Both the EO and the fungicide were diluted as described above and added separately to the culture medium. Petri plates were gently shaken before inoculation to enhance uniformity. All treatments were incubated in a culture chamber under the same conditions. After incubation, data expressed as *PGI* was calculated and assessed. Because of the great number of available methodologies, the interaction of the combination was determined using three methods: the additive method, the Abbott method, and the *FIC* index. The additive and Abbott methods have been traditionally used to determine interactions among agricultural pesticides (Kosman and Cohen 1996). At present, the *FIC* index is the most recommended method to evaluate combinations of antifungal compounds (Odds 2003). In the additive method, the expected *PGI* (*APGI*) for each mixture was calculated by adding the individual *PGI* of the fungicide doses (PGI_F) and the individual *PGI* of the EO doses (PGI_{EO}). Synergism, additive or antagonistic effects were determined when the observed *PGI* (*OPGI*) was statistically higher, equal and lower than the calculated *APGI*, respectively (Camiletti *et al.* 2016). In the Abbott method, the expected *PGI* (*EPGI*) was estimated also using the PGI_F and PGI_{EO} but following the equation:

$$EPGI = PGIF + PGI_{EO} - [(PGIF \times PGI_{EO})/100].$$

The effect of the combinations was designated by assessing the Abbot Index (*AI*):

$$AI = OPGI/EPGI.$$

A synergistic interaction was assigned for $AI \geq 1.5$, additive for $>0.5-1.4$ and antagonistic for <0.5 (Kosman and Cohen 1996).

For each combination, the utilized fungicide (C_{F+EO}) and EO (C_{EO+F}) concentrations served as a basis to calculate a fractional inhibitory concentration (*FIC*) index for the mixture according to the formulas:

$$FIC_{EO} = C_{EO+F}/C_{EO},$$

$$FIC_F = C_{F+EO}/C_F,$$

$$FIC = FIC_{EO} + FIC_F,$$

where: C_{EO} and C_F are the EOs and fungicide concentrations needed to achieve the same *OPGI* when they are applied individually. Predicted values of C_{EO} and C_F were obtained from regression equations. The interaction was interpreted as synergy for $FIC < 0.5$, additive for $>0.5-4.0$ and antagonistic for >4.0 (Pillai *et al.* 2005; Magi *et al.* 2015; Nikkhah *et al.* 2017).

Statistical analysis

Treatments were replicated three times and data of each experiment were combined after checking for homogeneity of the experimental error variances by the *F* test. Means were obtained via ANOVA ($p < 0.05$). Significant differences between treatments were determined using Fisher's protected least significant difference (LSD) test at level $p < 0.05$. The *PGI* data were regressed against the the log-transformed dose, and the best fit model was selected according to the significance of the regression and coefficient of determination (R^2). Regression equations were used to obtain the predicted values of the minimum inhibitory concentrations (*MICs*; $PGI = 50\%$), C_{EO} and C_F (Marei *et al.* 2012). Data analysis was performed using InfoStat software version 2014.

Results

Essential oil yield and chemical composition

The EO of *T. filifolia* was collected as a 4.0% dry weight yield and its composition was characterized by a great proportion of phenylpropanoids (Table 2). The main components were *cis*-anethole and estragole, representing 97.9% of the oil composition. *P*-Anisaldehyde, β -bisabolene, methyl eugenol and spatululenol completed the chemical profile with percentages lower than 1.0%.

Table 2. Chemical profile and relative percentage of main compounds present in the *Tagetes filifolia* essential oil according to GC-MS analysis

<i>R</i> ^a	<i>RT</i> ^b	Compound	Relative percentage [%]
1195	11.88	Estragole	22.11 ± 2.9
1252	12.55	p-Anisaldehyde	0.49 ± 0.3
1289	12.93	Anethole	75.10 ± 5.7
1401	13.89	Methyl eugenol	0.55 ± 0.2
1509	14.85	β-Bisabolene	0.69 ± 0.2
1576	15.59	Sphatulenol	0.26 ± 0.1
Total			99.20

^a*R* – retention index, ^b*RT* – retention time

Antifungal activity

The results of the antifungal activity screening are presented in Table 3. All the studied compounds (F1, F2, F3 and the EO) had the capacity to completely inhibit the growth of *C. truncatum* and *T. harzianum*. When the fungi were compared, *T. harzianum* showed a remarkably higher resistance than *C. truncatum* to all four antifungal agents. The fungicide F1 showed the strongest antifungal activity against both fungi, with the lowest being *MIC* and *MFC*, followed by F2 and F3. Surprisingly, the sensitivity of the pathogen to

F1 decreased at concentrations higher than the *MFC* (Fig. 1). The chemical fungicide F1 also affected the growth of *T. harzianum*, 10% on average, at the same doses that controlled *C. truncatum* (Fig. 1). The chemical fungicide F2 did not show an antifungal effect on *T. harzianum* when it was evaluated at the same concentration that controlled *C. truncatum*. In fact, the inhibitory effect against the biocontrol fungus started at a concentration of 10 μl · l⁻¹. The fungicide F3 presented the lowest antifungal activity among the chemical pesticides, showing the highest *MFC* against *C. truncatum*. Furthermore, growth inhibition of *T. harzianum* by this pesticide was not observed up to concentrations higher than 25 μl · l⁻¹. The sensitivity of *C. truncatum* to the EO was lower than to chemical fungicides. Furthermore, doses of EO that partially inhibited the growth of *C. truncatum* also affected mycelial development of *T. harzianum* (Fig. 2). For example, 4,000 μl · l⁻¹ of EO showed a *PGI* of 50 and 21% against *C. truncatum* and *T. harzianum*, respectively.

Mixture effects

The synergism assay was carried out to find effective combinations of fungicides and EO that allow a reduction of the pesticide dose required to control *C. truncatum*. The objective was to find combinations

Table 3. Minimum fungicidal concentration of chemical compounds and *Tagetes filifolia* essential oil against *Colletotrichum truncatum* and *Trichoderma harzianum*

Fungi	Essential oil ^a		Fungicides ^b					
	<i>Tagetes filifolia</i>		F1		F2		F3	
	<i>MIC</i> ^c	<i>MFC</i>	<i>MIC</i>	<i>MFC</i>	<i>MIC</i>	<i>MFC</i>	<i>MIC</i>	<i>MFC</i>
<i>Colletotrichum truncatum</i>	4,000	6,000	0.13	0.25	0.35	1.0	0.7	1.5
<i>Trichoderma harzianum</i>	7,400	9,000	1.10	1.50	10.00	30.0	75.0	90.0

^amean values (*n* = 3) expressed as μl · l⁻¹ of essential oil

^bF1 – carbendazim, F2 – difenconazole, F3 – trifloxystrobin + cyproconazole. Mean values (*n* = 3) expressed as μl · l⁻¹ of active ingredient

^c*MIC* – minimum inhibitory concentration, *MFC* – minimum fungicidal concentration




Fig. 1. Percentage of growth inhibition (*PGI*) of *Colletotrichum truncatum* and *Trichoderma harzianum* exposed to different concentrations of carbendazim




Fig. 2. Percentage of growth inhibition (*PGI*) of *Colletotrichum truncatum* and *Trichoderma harzianum* exposed to different concentrations of *Tagetes filifolia* EO

Table 4. Antifungal activity of chemical fungicides combined with *Tagetes filifolia* essential oil against *Colletotrichum truncatum* and their interactions according to the additive, Abbott and FIC methods

Compound A		Compound B		PGI		Additive effect	Abbott		FIC	
fungicide ^a	concentration ^b	EO concentration ^b	observed	expected	index		effect	index	effect	
F1	20	20	11 a	4 b	SN	2.89	SN	0,84	AD	
F1	20	40	31 a	4 b	SN	8.00	SN	1.03	AD	
F1	20	60	59 a	27 b	SN	2.22	SN	1.19	AD	
F1	20	80	94 a	85 a	AD	1.11	AD	1.11	AD	
F1	40	20	64 a	4 b	SN	16.61	SN	0.91	AD	
F1	40	40	73 a	4 b	SN	18.62	SN	1.10	AD	
F1	40	60	75 a	27 b	SN	2.80	SN	1.34	AD	
F1	40	80	90 a	86 a	AD	1.06	AD	1.35	AD	
F1	60	20	78 a	62 a	AD	1.29	AD	0.99	AD	
F1	60	40	84 a	62 b	SN	1.39	AD	1.14	AD	
F1	60	60	100 a	85 b	SN	1.44	AD	1.20	AD	
F1	60	80	100 a	100 b	AD	1.07	AD	1.40	AD	
F1	80	20	84 a	62 b	SN	1.40	AD	1.09	AD	
F1	80	40	93 a	62 b	SN	1.55	SN	1.26	AD	
F1	80	60	96 a	85 b	SN	1.38	AD	1.47	AD	
F1	80	80	100 a	100 a	AD	1.07	AD	1.60	AD	
F2	20	20	20 b	37 a	AN	0.58	AD	1.79	AD	
F2	20	40	25 a	34 a	AD	0.77	AD	1.74	AD	
F2	20	60	61 a	57 a	AD	1.25	AD	1.29	AD	
F2	20	80	100 a	100 a	AD	1.09	AD	1.08	AD	
F2	40	20	1 b	70 a	AN	0.00	AN	22.00	AN	
F2	40	40	10 b	67 a	AN	0.16	AN	20.83	AN	
F2	40	60	69 b	91 a	AN	0.94	AD	1.60	AD	
F2	40	80	100 a	100 a	AD	1.03	AD	1.27	AD	
F2	60	20	28 b	77 a	AN	0.40	AN	3.07	AD	
F2	60	40	31 b	74 a	AN	0.44	AN	3.17	AD	
F2	60	60	53 b	98 a	AN	0.69	AD	2.42	AD	
F2	60	80	100 a	100 a	AD	1.05	AD	1.40	AD	
F2	80	20	30 b	92 a	AN	0.35	AN	3.81	AD	
F2	80	40	33 b	90 a	AN	0.39	AN	3.86	AD	
F2	80	60	47 b	100 a	AN	0.53	AD	3.19	AD	
F2	80	80	100 a	100 a	AD	1.03	AD	1.60	AD	
F3	20	20	35 a	17 b	SN	2.15	SN	0.56	AD	
F3	20	40	53 a	10 b	SN	5.43	SN	0.75	AD	
F3	20	60	67 a	19 b	SN	3.72	SN	0.96	AD	
F3	20	80	88 a	69 b	SN	1.40	AD	1.09	AD	
F3	40	20	36 a	11 b	SN	3.31	SN	0.84	AD	
F3	40	40	54 a	4 b	SN	14.46	SN	1.01	AD	
F3	40	60	77 a	13 b	SN	6.13	SN	1.14	AD	
F3	40	80	93 a	62 b	SN	1.54	AD	1.26	AD	
F3	60	20	36 a	28 a	AD	1.35	AD	1.12	AD	
F3	60	40	73 a	21 b	SN	3.42	SN	1.15	AD	
F3	60	60	82 a	30 b	SN	2.87	SN	1.33	AD	
F3	60	80	100 a	80 b	SN	1.41	AD	1.45	AD	
F3	80	20	78 a	60 b	SN	1.40	AD	1.11	AD	
F3	80	40	86 a	52 b	SN	1.64	AD	1.29	AD	
F3	80	60	90 a	61 b	SN	1.59	AD	1.49	AD	
F3	80	80	100 a	100 a	AD	1.24	AD	1.60	AD	

PGI – percentage of growth inhibition; FIC – fractional inhibitory concentration

^aF1 – carbendazim, F2 – difenoconazole, F3 – trifloxystrobin (70%) + cyproconazole (30%)^bconcentration expressed as percentages of their minimum fungicidal concentrations.The same letter in the row means that there are no significant differences at $p < 0.05$ ($n = 3$; Fisher's LSD test); SN – synergism, AD – additivity, AN – antagonism

of fungicides with *T. filifolia* EO that improved their performance and to study their effects on the growth of the biocontrol agent *T. harzianum*. A total of 48 combinations were evaluated: 20, 40, 60 or 80% of the MFC of each fungicide combined with 20, 40, 60 or 80% of the MFC of EO. Synergistic, additive or antagonistic effects were assigned based on three different methods: the additive method, the Abbott method and the FIC index (Table 4). Synergistic interactions were not found according to the FIC index, being mainly determined as an additive type. The Abbott and additive methods displayed some synergistic interactions although none of them inhibited completely the growth of *C. truncatum*. However, the EO presented a total of eight combinations with different fungicides that allowed reductions of the pesticide doses, conserving its antifungal effect. Combinations of EO with F2 showed the largest number of additive combinations controlling *C. truncatum*. In fact, the mixture of 80% EO MFC and 20% F2 MFC showed a 100% inhibition which allowed an 80% reduction of the pesticide dose. However, these antifungal agents also presented a greater number of antagonist interactions when they were combined. On the other hand, all combinations that allowed a reduction of the pesticide dose against *C. truncatum* affected the growth of *T. harzianum*. The PGIs averaged $31 \pm 4\%$, in accordance with the PGIs observed when 60 and 80% of EO MFC were evaluated individually (Fig. 2).

Discussion

In Argentina, soybean crops are affected by anthracnose, causing yield and quality losses. Soybean monocropping and conservation tillage have exacerbated the severity of anthracnose, increasing the doses of fungicides applied to the crop (Carmona et al. 2015). The *T. filifolia* EO was characterized and investigated as a natural alternative to reduce the amount of chemical pesticides currently applied to soybean crops.

The EO yield obtained from the *T. filifolia* aerial parts was lower than those reported for plants grown in other regions of Argentina (6.3–13.1%), Mexico (6.6–16.8%) and Peru (12.1%) (Maestri et al. 1991; Serrato Cruz et al. 2003; Marotti et al. 2011). Although the obtained EO yield was lower than that reported in earlier literature, the chemical composition showed a similar profile to those previously reported for *T. filifolia* EO. The prevalence of *cis*-anethole and estragole was reported in EO extracted from different cultivars grown in other regions of Argentina, in agreement with our results (Maestri et al. 1991; Zygadlo et al. 1994; Camiletti et al. 2016). In addition, the EO obtained

from aerial parts of *T. filifolia* grown in Peru was previously characterized by a similar composition (De Feo et al. 1998). Studies by Maestri et al. (1991) indicated that EO yield is related to the altitude above sea level of the plant population, acting as an adaptive response mechanism that does not influence the chemical composition of the essence.

All the studied compounds showed antifungal activity against the pathogen *C. truncatum* and the biocontrol agent *T. harzianum*. Among the chemical fungicides, F1 showed the strongest antifungal activity, followed by F2 and F3. This finding is in agreement with other research where members of the benzimidazole group needed lower concentrations to control *C. truncatum* than members of other chemical groups (Zavala León et al. 2005). The effectiveness of F1 to control *C. truncatum* isolates was previously demonstrated in studies that included several strains of this pathogen isolated from other crops and regions (Gawade et al. 2009; Dias et al. 2016). Moreover, Oh and Kang (2002) observed that the antifungal activity of this fungicide decreased at concentrations higher than those that completely controlled *C. truncatum*, which agrees with results obtained in this work. The active ingredients difenoconazole (F2) and cyproconazole (present in F3) were described by a similar antifungal activity against *Colletotrichum* species (Chen et al. 2016). Cyproconazole represented only 30% of the active ingredient in F3 and trifloxystrobin (strobilurin), the remaining 70%. In previous studies (Carmona et al. 2011; Rampersad et al. 2012; Torres-Calzada et al. 2015), *C. truncatum* isolates showed intermediate or higher resistance to fungicide members of the chemical group strobilurins, explaining the lower inhibitory effect of F3. In fact, F2 and F3 did not affect the growth of *T. harzianum* at the same concentration that controlled the pathogen as was observed for F1. This biocontrol agent was investigated by its sensitivity to numerous fungicides, indicating that diverse fungicides can inhibit the growth of this fungus (Sarkar et al. 2010; Sushir et al. 2015).

Tagetes filifolia EO was effective in controlling *C. truncatum*. This EO was previously described by its antifungal activity against several fungi, including species of genera *Alternaria*, *Colletotrichum* and *Sclerotium* (Zygadlo 1994; Camiletti et al. 2016). The antimicrobial activity of the major compound in the EO of *T. filifolia* explained the antifungal property of this essence. Anethole demonstrated an inhibitory effect against species of genera *Alternaria*, *Aspergillus*, *Fusarium* and *Rhizoctonia*, among others (Karapinar 1990; Huang et al. 2010). The efficacy of the *T. filifolia* EO against *C. truncatum* was lower than that observed for the chemical fungicides, which is in agreement with previous research (Camiletti et al. 2016). Furthermore,

this EO affected mycelial development of *T. harzianum*. In fact, some EOs were reported as effective against *T. harzianum* in crops where this fungus is considered to be a plant pathogen (Angelini *et al.* 2008). These results differ from those reported in earlier studies where it was indicated that *T. harzianum* improved the performance of EOs against several phytopathogenic fungi in combined treatments (Abdel-Kader *et al.* 2011). However, the effectiveness of the biocontrol agent by itself was not included in the study carried out by Abdel-Kader *et al.* (2011).

Combinations of fungicides with *T. filifolia* EO showed the prevalence of additive interactions. A similar characterization was previously indicated for several EOs that exhibited this type of interaction when combined with chemical fungicides, such as fludioxonil and iprodione (Camiletti *et al.* 2016; Camiletti 2018). Strong synergistic interactions have been mainly documented for mixtures of antifungal agents with different modes of action (Gisi *et al.* 1985; Gisi 1991, 1996; Kosman and Cohen 1996; Camiletti *et al.* 2016). Carbendazim inhibits the polymerization of tubulin, a protein that is essential during cell division (FAO 2005). Triazoles difenoconazole and cyproconazole act by inhibition of ergosterol biosynthesis, affecting the structure and function of cell membranes (Szkolnik 1981). Trifloxystrobin works by interfering with respiration and its site of action is located in the mitochondrial respiratory pathway (FAO 2005). The terpenes commonly present in EOs were previously described as damaging factors of membranous structures, depleting hyphal cytoplasm and organelles (Lucini 2004; Bakkali *et al.* 2008; Plodpai *et al.* 2013). Triazoles and EOs have a similar mode of action and may explain the higher number of interactions with antagonistic effects. Despite different modes of action, combinations of carbendazim and trifloxystrobin with the EO did not display synergistic effects that controlled *C. truncatum*. However, these combinations were characterized by their additive interaction that remarkably reduced the fungicide dose. Thus, these mixtures could be used as a part of integrated management to reduce the ecological impact of chemical pesticides (Thompson and Kreutrweiser 2007). These properties were widely investigated in combinations of EOs with medicinal fungicides (Shin and Kang 2003; Pyun and Shin 2006; Amber *et al.* 2010). Some research has been carried out to investigate combinations of EOs with agricultural fungicides. Some EOs were reported as an alternative to reduce the amount of the chemical fungicide fludioxonil needed to control *A. flavus* and *Penicillium* spp. (Camiletti 2018). The *T. filifolia* EO was reported as a natural antifungal agent against *Sclerotium cepivorum* with a remarkable capacity to reduce the required dose of the fungicide iprodione (Camiletti *et al.* 2016).

The inclusion of natural compounds as part of an integrated management program is critical for controlling plant diseases. This study demonstrates that the antifungal power of the *T. filifolia* EO against *C. truncatum* can be exploited to reduce chemical doses of fungicides commonly applied to soybean crops. *T. filifolia* EO has the potential to be applied individually as well as in combination with chemical fungicides. The antifungal effect of this EO against *T. harzianum* must be considered in the development of new strategies that may include this biocontrol strain. In conclusion, *T. filifolia* EO is a novel alternative to control anthracnose caused by *C. truncatum* and can be used to reduce the amount of chemical fungicide applied to the field. The agrochemical industry should consider *T. filifolia* EO and its main components for the production of environmentally friendly fungicides. Moreover, the persistence of fungicide residues and the amount of money spent on them can be diminished. Further field assays are needed to adjust the applied concentrations.

Acknowledgements

The authors wish to thank the National Institute of Agricultural Technology (INTA), National University of Cordoba, Secretary of Science and Technology (UNC, SECyT) and the Argentinian National Research Council (CONICET) for their financial support. We are also grateful to the College of Agricultural Science (FCA-UNC) for allowing us to use its laboratory equipment.

References

- Abbaszadeh S., Sharifzadeh A., Shokri H., Khosravi A.R., Abbaszadeh A. 2014. Antifungal efficacy of thymol, carvacrol, eugenol and menthol as alternative agents to control the growth of food-relevant fungi. *Journal of Mycologie Medicale* 24 (2): e51–e56. DOI: <https://doi.org/10.1016/j.mycmed.2014.01.063>
- Abdel-Kader M., El-Mougy N., Lashin S. 2011. Essential oils and *Trichoderma Harzianum* as an integrated control measure against faba bean root rot pathogens. *Journal of Plant Protection Research* 51 (3): 306–313. DOI: <https://doi.org/10.2478/v10045-011-0050-8>
- Abrol D.P., Shankar U. 2016. Integrated Pest Management. p. 523–549. In: “Breeding Oilseed Crops for Sustainable Production” (S.K. Gupta, ed.). Elsevier Inc. Netherlands, 994 pp.
- Adams R.P. 1995. Identification of Essential Oils by Ion Trap Mass Spectroscopy. Carol Stream, IL, USA, Allured Publishing, 843 pp.
- Amber K., Aijaz A., Immaculata X., Luqman K.A., Nikhat M. 2010. Anticandidal effect of *Ocimum sanctum* essential oil and its synergy with fluconazole and ketoconazole. *Phyto-medicine* 17: 921–925. DOI: [10.1016/j.phymed.2010.02.012](https://doi.org/10.1016/j.phymed.2010.02.012)
- Angelini P., Pagiotti R., Granetti B. 2008. Effect of antimicrobial activity of *Melaleuca alternifolia* essential oil on antagonistic potential of *Pleurotus* species against *Trichoderma*

- harzianum* in dual culture. World Journal of Microbiology and Biotechnology 24: 197–202. DOI: 10.1007/s11274-007-9456-x
- Armstrong-Cho C.L., Banniza S. 2006. *Glomerella truncata* sp. nov., the teleomorph of *Colletotrichum truncatum*. Mycological Research 110 (8): 951–956. DOI: <https://doi.org/10.1016/j.mycres.2006.06.002>
- Arrebola E., Sivakumar D., Bacigalupo R., Korsten L. 2010. Combined application of antagonist *Bacillus amyloliquefaciens* and essential oils for the control of peach postharvest diseases. Crop Protection 29 (4): 369–377. DOI: <https://doi.org/10.1016/j.cropro.2009.08.001>
- Asensio C.M., Nepote V., Grosso N.R. 2011. Chemical stability of extra-virgin olive oil added with oregano essential oil. Journal of Food Science 76: 445–450. DOI: 10.1111/j.1750-3841.2011.02332.x
- Bakkali F., Averbeck S., Averbeck D., Idaomar M. 2008. Biological effects of essential oils – a review. Food Chemical Toxicology 46: 446–75. DOI: 10.1016/j.fct.2007.09.106
- Benítez T., Rincón A.M., Limón M.C., Codón A.C. 2004. Biocontrol mechanisms of *Trichoderma* strains. International Microbiology 7: 249–260.
- Bluma R., Amaldien M.R., Daghero J., Etcheverry M. 2008. Control of *Aspergillus* section *Flavi* growth and aflatoxin accumulation by plant essential oils. Journal of Applied Microbiology 105 (1): 203–214. DOI: <https://doi.org/10.1111/j.1365-2672.2008.03741.x>
- Camiletti B.X. 2018. Estrategias de manejo de *Aspergillus flavus* y *Penicillium* spp. para la reducción de los niveles de micotoxinas en maíz. Ph.D. Thesis, Universidad Nacional de Córdoba, Spain.
- Camiletti B.X., Asensio C.M., Gadban L.C., Pecci M. de la P.G., Conles M.Y., Lucini E.I. 2016. Essential oils and their combinations with iprodione fungicide as potential antifungal agents against the rot (*Sclerotium cepivorum* Berk) in garlic (*Allium sativum* L.) crops. Industrial Crops and Products 85: 117–124. DOI: <https://doi.org/10.1016/j.indcrop.2016.02.053>
- Camiletti B.X., Asensio C.M., Pecci M.D.L.P.G., Lucini E.I. 2014. Natural control of corn postharvest fungi *Aspergillus flavus* and *Penicillium* spp. using essential oils from plants grown in Argentina. Journal of Food Science 79 (12): M2499–M2506. DOI: <https://doi.org/10.1111/1750-3841.12700>
- Carmona M., Gally M., Sautua F., Abello A., Lopez P. 2011. Uso de mezclas de azoxistrobina y triazoles para controlar enfermedades de fin de ciclo de la soja. [Use of azoxystrobin and triazoles mixtures to control late season diseases in soybean crop]. Summa Phytopathologica 37 (2): 134–139. DOI: <https://doi.org/10.1590/s0100-54052011000200009> (in Spanish, with English summary)
- Carmona M., Sautua F., Perelman S., Gally M., Reis E.M. 2015. Development and validation of a fungicide scoring system for management of late season soybean diseases in Argentina. Crop Protection 70: 83–91. DOI: 10.1016/j.cropro.2015.01.019
- Chen S.N., Luo C.X., Hu M.J., Schnabel G. 2016. Sensitivity of *Colletotrichum* species, including *C. fioriniae* and *C. nymphaeae*, from peach to demethylation inhibitor fungicides. Plant Disease 100: 2434–2441. DOI: 10.1094/PDIS-04-16-0574-RE
- Dhingra O.D., Sinclair J.B. 1995. Basic Plant Pathology Methods. CRC Press, Florida, United States, 444 pp.
- Dias M.D., Pinheiro V.F., Café-filho A.C. 2016. Impact of anthracnose on the yield of soybean subjected to chemical control in the north region of Brazil. Summa Phytopathologica 42 (1): 18–23. DOI: <https://doi.org/10.1590/0100-5405/2114>
- FAO. 2005. Pesticide residues in food. Available on: http://www.fao.org/fileadmin/templates/agphome/documents/Pests_Pesticides/JMPR/JMPR05report.pdf [Accessed: May 25, 2019]
- De Feo V., Della Porta G., Urrunaga Soria E., Urrunaga Soria R., Senatore F. 1998. Composition of the essential oil of *Tagetes filifolia* Lag. Flavour Fragrance Journal 13: 145–147. DOI: 10.1002/(SICI)1099-1026(199805/06)
- Fernandez-Ortuño D., Chen F., Schnabel G. 2012. Resistance to pyraclostrobin and boscalid in *Botrytis cinerea* isolates from strawberry fields in the Carolinas. Plant Disease 96 (8): 1198–1203. DOI: <https://doi.org/10.1094/pdis-12-11-1049-re>
- Ganuza M., Pastor N., Erazo J., Andrés J., Reynoso M.M., Rovera M., Torres A.M. 2017. Efficacy of the biocontrol agent *Trichoderma harzianum* ITEM 3636 against peanut smut, an emergent disease caused by *Thecaphora frezii*. European Journal of Plant Pathology 151 (1): 257–262. DOI: <https://doi.org/10.1007/s10658-017-1360-0>
- Gawade D., Phule M., Vidyapeeth K., Gawade D. 2009. Field evaluation of fungicides, botanicals and bioagents against anthracnose of soybean. Agricultural Science 29: 174–177.
- Gisi U. 1991. Synergism between fungicides for control of *Phytophthora*. p. 361–372. In: “Phytophthora” (J.A. Lucas, D.S. Shattock, L.R. Cooke, eds.). Cambridge University Press, Cambridge, UK, 1089 pp.
- Gisi U. 1996. Synergistic interactions of fungicides in mixtures. Phytopathology 86: 1273–1279.
- Gisi U., Binde H., Rimbach E. 1985. Synergistic interactions of fungicides with different modes of action. Transactions of the British Mycological Society 85: 299–306. DOI: 10.1016/S0007-1536(85)80192-3
- Goncalves S.S., Souza A.C.R., Chowdhary A., Meis J.F., Colombo A.C. 2016. Epidemiology and molecular mechanisms of antifungal resistance in *Candida* and *Aspergillus*. Mycoses 59: 198–219. DOI: 10.1111/myc.12469
- Grondona I., Hermosa R., Tejada M., Gomis M.D., Mateos P.F., Bridge P.D. 1997. Physiological and biochemical characterization of *Trichoderma harzianum*, a biological control agent against soilborne fungal plant pathogens. Applied Environmental Microbiology 63: 3189–3198. DOI: 0099-2240/97/\$04.0010
- Huang Y., Zhao J., Zhou L., Wang J., Gong Y., Chen X. 2010. Antifungal activity of the essential oil of *Illicium verum* fruit and its main component trans-anethole. Molecules 15: 7558–7569. DOI: 10.3390/molecules15117558
- Isman M.B. 2000. Plant essential oils for pest and disease management. Crop Protection 19 (8–10): 603–608. DOI: [https://doi.org/10.1016/s0261-2194\(00\)00079-x](https://doi.org/10.1016/s0261-2194(00)00079-x)
- Karapinar M. 1990. Inhibitory effects of anethole and eugenol on the growth and toxin production of *Aspergillus parasiticus*. International Journal of Food Microbiology 10: 193–199. DOI: 10.1016/0168-1605(90)90066-E
- Kosman E., Cohen Y. 1996. Procedures for calculating and differentiating synergism and antagonism in action of fungicide mixtures. Phytopathology 86: 1263–1272.
- Lim S., Shin S. 2008. Combined effects of the essential oil from *Eucalyptus globulus* with ketoconazole against *Candida* and *Trichophyton* species. Natural Products Science 14 (1): 1–12.
- Lucini E.I. 2004. Cambios en la composición lipídica del micelio de *Sclerotinia sclerotiorum* (Lib.) de Bary y *Sclerotium cepivorum* Berk por acción de terpenos. Ph.D. Thesis. Universidad Nacional de Córdoba, Spain.
- Lucini E.I., Zunino M.P., Lopez M.L., Zygadlo J.A. 2006. Effect of monoterpenes on lipid composition and sclerotial development of *Sclerotium cepivorum* Berk. Journal of Phytopathology 154 (7–8): 441–446. DOI: <https://doi.org/10.1111/j.1439-0434.2006.01126.x>
- Maestri D.M., Zygadlo J.A., Grosso N.R., Aburrá R.E. 1991. Composición del aceite esencial en poblaciones de *Tagetes filifolia*. Annals of Biology 17: 133–136.
- Magi G., Marini E., Facinelli B. 2015. Antimicrobial activity of essential oils and carvacrol, and synergy of carvacrol and erythromycin against clinical erythromycin-resistant Group A Streptococci. Frontiers in Microbiology 6: 1–7. DOI: <https://doi.org/10.3389/fmicb.2015.00165>
- Manandhar J.B., Hartman G.L. 1999. Anthracnose. p. 13–14. In: “Compendium of Soybean Disease” (G.L. Hartman,

- J.B. Sinclair, J.C. Rupe, eds.). American Phytopathological Society, St. Paul, MN, 238 pp.
- Marei G.I.K., Abdel Rasoul M.A., Abdelgaleil S.A.M. 2012. Comparative antifungal activities and biochemical effects of monoterpenes on plant pathogenic fungi. *Pesticide Biochemistry and Physiology* 103 (1): 56–61. DOI: <https://doi.org/10.1016/j.pestbp.2012.03.004>
- Marotti M., Piccaglia R., Biavati B., Marotti I., Marotti M., Piccaglia R. 2011. Characterization and yield evaluation of essential oils from different tagetes species. *Journal of Essential Oil Research* 15: 37–41. DOI: <https://doi.org/10.1080/10412905.2004.9698767>
- Nikkhah M., Hashemi M., Habibi Najafi M.B., Farhoosh R. 2017. Synergistic effects of some essential oils against fungal spoilage on pear fruit. *International Journal of Food Microbiology* 257: 285–294. DOI: <https://doi.org/10.1016/j.ijfoodmicro.2017.06.021>
- Odds F.C. 2003. Synergy, antagonism, and what the checkerboard puts between them. *Journal of Antimicrobial Chemotherapy* 52 (1): 1. DOI: <https://doi.org/10.1093/jac/dkg301>
- Oh J.H., Kang N.W. 2002. Efficacy of fluazinam and Iprodione + Propineb in the suppression of *Diaporthe phaseolorum*, *Colletotrichum truncatum* and *Cercospora kikuchii*, the causal agents of seed decay in soybean. *Plant Pathology Journal* 18 (4): 216–220. DOI: <https://doi.org/10.5423/ppj.2002.18.4.216>
- Omidbeygi M., Barzegar M., Hamidi Z., Naghdibadi H. 2007. Antifungal activity of thyme, summer savory and clove essential oils against *Aspergillus flavus* in liquid medium and tomato paste. *Food Control* 18 (12): 1518–1523. DOI: <https://doi.org/10.1016/j.foodcont.2006.12.003>
- Pavela R. 2014. Acute, synergistic and antagonistic effects of some aromatic compounds on the *Spodoptera littoralis* Bois. (Lep., Noctuidae) larvae. *Industrial Crops and Products* 60: 247–258. DOI: <https://doi.org/10.1016/j.indcrop.2014.06.030>
- Phélinas P., Choumert J. 2017. Is GM soybean cultivation in Argentina sustainable? *World Development* 99: 452–462. DOI: [10.1016/j.worlddev.2017.05.033](https://doi.org/10.1016/j.worlddev.2017.05.033)
- Pillai S.K., Moellering R.C., Eliopoulos G.M. 2005. Antimicrobial combinations. p. 365–340. In: “Antibiotics in Laboratory Medicine” (V. Lorian, ed.). Lippincott, Williams and Wilkin, Philadelphia, USA, 1099 pp.
- Plodpai P., Chuenchitt S., Petcharat V., Chakthong S., Voravuthikunchai S.P. 2013. Anti-*Rhizoctonia solani* activity by *Desmos chinensis* extracts and its mechanism of action. *Crop Protection* 43: 65–71. DOI: [10.1016/j.cropro.2012.09.004](https://doi.org/10.1016/j.cropro.2012.09.004)
- Pyun M.S., Shin S. 2006. Antifungal effects of the volatile oils from *Allium* plants against *Trichophyton* species and synergism of the oils with ketoconazole. *Phytomedicine* 13 (6): 394–400. DOI: <https://doi.org/10.1016/j.phymed.2005.03.011>
- Ramos A.M., Tadic F.L., Cinto I., Carmona M., Gally M. 2013. Molecular characterization of *Colletotrichum* species causing soybean anthracnose in Argentina. *Mycotaxon* 123 (1): 457–465.
- Rampersad S.N., Teelucksingh L.D., Indies W., Sciences L., Science F. 2012. Differential responses of *Colletotrichum gloeosporioides* and *C. truncatum* isolates from different hosts to multiple fungicides based on two assays. *Plant Disease* 96 (10): 1526–1536. DOI: <https://doi.org/10.1094/pdis-10-11-0906-re>
- Rifai M.A. 1969. A revision of the genus *Trichoderma*. *Mycological Papers* 116: 1–56.
- Sarkar S., Narayanan P., Divakaran A., Balamurugan A., Premkumar R. 2010. The *in vitro* effect of certain fungicides, insecticides, and biopesticides on mycelial growth in the biocontrol fungus *Trichoderma harzianum*. *Turkish Journal of Biology* 34: 399–403. DOI: [10.3906/biy-0812-4](https://doi.org/10.3906/biy-0812-4)
- Serrato Cruz M.A., Reyes Trejo B., Ortega Arenas L., Domingo Gómez A., Gómez Sánchez N., López Martínez F. 2003. Anisillo (*Tagetes filifolia* Lag.): Recurso genético mexicano para controlar la mosquita blanca (*Bemisia* sp. y *Trialurodes* sp.). *Revista del Jardín Botánico Nacional* 24: 65–70.
- Sharma V., Salwan R., Sharma P.N., Kanwar S.S. 2017. Elucidation of biocontrol mechanisms of *Trichoderma harzianum* against different plant fungal pathogens: Universal yet host specific response. *International Journal of Biological Macromolecules* 95: 72–79. DOI: <https://doi.org/10.1016/j.ijbiomac.2016.11.042>
- Shin S., Kang C.A. 2003. Antifungal activity of the essential oil of *Agastache rugosa* Kuntze and its synergism with ketoconazole. *Letters in Applied Microbiology* 36 (2): 111–115. DOI: <https://doi.org/10.1046/j.1472-765x.2003.01271.x>
- Sosa-Gomez D.R., Delpin K.E., Moscardi F., Nozaki M. 2003. The impact of fungicides on *Nomurea rileyi* (Farlow) Samson epizootics and populations of *Anticarsia gemmatalis* Hübner (Lepidoptera: Noctuidae), on soybean. *Neotropical Entomology* 32 (2): 287–291. DOI: <https://doi.org/10.1590/s1519-566x2003000200014>
- Sushir M.A., Suryawanshi K.K., Patole S.P. 2015. Sensitivity of *Trichoderma harzianum* Rifai against systemic fungicides. *International Journal of Applied Research* 1: 403–405.
- Szkolnik M. 1981. Physical modes of action of sterol-inhibiting fungicides against apple diseases *Venturia inaequalis*. *Plant Disease* 65 (12): 981–985. DOI: <https://doi.org/10.1094/pd-65-981>
- Thompson D.G., Kreutweiser D.P. 2007. A review of the environmental fate and effects of natural “reduced-risk” pesticides in Canada. *ACS Symposium* 947: 245–274.
- Torres-Calzada C., Tapia-Tussell R., Higuera-Ciajara I., Martin-Mex R., Nexticapan-Garcez A., Perez-Brito D. 2015. Sensitivity of *Colletotrichum truncatum* to four fungicides and characterization of thiabendazole-resistant isolates. *Plant Disease* 99 (11): 1590–1595. DOI: <https://doi.org/10.1094/pdis-11-14-1183-re>
- Vollmann J. 2016. Soybean versus other food grain legumes: A critical appraisal of the United Nations International Year of Pulses 2016. *Journal of Land Management, Food Environment* 67: 17–24. DOI: [10.1515/boku-2016-0002](https://doi.org/10.1515/boku-2016-0002)
- Zavala León M.J., Tun Suárez J.M., Cristóbal Alejo J., Ruiz Sánchez E., Gutierrez Alonso O., Vázquez Calderón M. 2005. Control postcosecha de la antracnosis en papaya y sensibilidad de *Colletotrichum gloeosporioides* (Penz.) Sacc. a fungicidas organosintéticos. *Revista Chapingo – Serie Horticultura* 11: 251–255.
- Zygadlo J., Guzman C.A., Grosso N.R. 1994. Antifungal properties of the leaf oils of *Tagetes minuta* L. and *Tagetes filifolia* Lag. *Journal of Essential Oil Research* 6: 617–621. DOI: [10.1080/10412905.1994.9699353](https://doi.org/10.1080/10412905.1994.9699353)