

Towards a standard Tithonian to Valanginian calpionellid zonation of the Tethyan Realm

ISKRA LAKOVA¹ AND SILVIYA PETROVA²

Geological Institute, Bulgarian Academy of Sciences, Acad. G. Bonchev St., Bl. 24, 1113 Sofia, Bulgaria.

¹E-mails: lakova@geology.bas.bg, ²silviya_p@geology.bas.bg

ABSTRACT:

Lakova, I. and Petrova, S. 2013. Towards a standard Tithonian to Valanginian calpionellid zonation of the Tethyan Realm. *Acta Geologica Polonica*, **63** (2), 201–221. Warszawa.

A calpionellid zonal scheme is proposed for the Tithonian through Valanginian pelagic carbonates of the Western Balkan Unit, based on the vertical ranges of 57 chitinoideid and calpionellid species recognized. This zonal scheme consists of calpionellid zones that are widely accepted in the Mediterranean Realm, such as the zones of *Chitinoidea*, *Praetintinnopsella*, *Crassicollaria*, *Calpionella*, *Calpionellopsis*, *Calpionellites* and *Tintinnopsella*. Subzonal divisions are comparable to those in the Carpathians. Direct correlations between ammonite and calpionellid ranges suggest that the base of the Upper Tithonian corresponds to the FO of *Chitinoidea boneti*; that of the Upper Berriasian to the FO of the genus *Calpionellopsis*; and the base and top of the Lower Valanginian to the FO and LO of the genus *Calpionellites* respectively. Correlations of the calpionellid zonation in the study area with zonations in other areas are discussed.

Key words: Calpionellids; Tithonian; Berriasian; Valanginian; Western Balkan Unit; Mediterranean standard zonation.

INTRODUCTION

The pelagic carbonate sequences cropping out in the West Balkan Mountains, western Bulgaria (Text-fig. 1) provide an excellent calpionellid record through the Jurassic–Cretaceous boundary interval. The boundary succession, spanning the Tithonian, Berriasian and Valanginian, is represented lithostratigraphically by the Gintsi Formation (pink and grey nodular limestones), Glozhene Formation (hard grey micritic limestones) and the Salash Formation (alternation of micritic limestones, clayey limestones and marls).

The lithostratigraphy and ammonite biostratigraphy of the successions studied were documented earlier (Nikolov and Sapunov 1970, 1977; Nikolov and Tsankov 1971; Mandov 1971; Sapunov 1976). The Gintsi and Glozhene formations were introduced by

Nikolov and Sapunov (1970), and the Salash Formation by Nikolov and Tsankov (1971). Mandov (1971) analysed the lithological sequences and ammonite faunas of the Lower Cretaceous parts of the successions.

The calpionellid biostratigraphy of the Tithonian to Valanginian of the Western Balkan Mts and the Western Fore-Balkan was established by Bakalova (1977), Lakova (1993, 1994), Lakova *et al.* (1999, 2007) and Ivanova *et al.* (2006). Bakalova (1977) recognized the zones of *Chitinoidea* through to *Calpionellites*, with two subzones (Alpina and Elliptica-Carpathica) distinguished in the *Calpionella* Zone. Bakalova-Ivanova (1986) divided the *Calpionella* Zone into the Alpina, Remaniella and Elliptica subzones.

Three main periods can be distinguished in the calpionellid biostratigraphic studies of the uppermost Jurassic and Lower Cretaceous. The first period, initiated by

Remane (1963, 1971), Allemann *et al.* (1971), Trejo (1980) and Borza (1984), ended with the establishment of the standard zonation for the Tethyan Realm. During the second period (1986–1999) the calpionellid zonation was refined, and the key bioevents and subzones were defined (Pop 1986c, 1989, 1994, 1997b; Bakalova-Ivanova 1986; Remane *et al.* 1986; Reháková 1995; Reháková and Michalík 1997a, b; Grün and Blau 1997; Lakova *et al.* 1999). The third period represents a renaissance of calpionellid research, with new projects spanning wider geographic territories, refining the taxonomy and biostratigraphy of the group, and integrating calpionellid studies with other biostratigraphies, as well as with chemo- and magnetostratigraphy.

The present paper aims at presenting the results of a calpionellid zonation for the Western Balkan Mts, western Bulgaria, with discussion of the correlations with all of the important previous zonations in the Tethyan Realm. The widely accepted calpionellid bioevents and subzones, and selected unresolved problems are discussed.

The calpionellid biostratigraphy of the Gintsi 1 and Gintsi 2 sections studied herein was the subject of SP's unpublished Ph.D. thesis "Ammonite and calpionellid biostratigraphy of the Berriasian, Valanginian and Hauterivian in the Western Srednogorie and Western Balkan Mts", supervised by IL. The calpionellids from the Barlya section have been re-studied using material from Lakova *et al.* (1999), reflecting more recent taxonomic concepts and knowledge of calpionellid bioevents.

The Barlya section is located immediately north of the village of Barlya, Sofia District (Text-fig. 2). A total of 57 thin sections were examined from this locality, spanning the top of the Gintsi Formation (samples 0310–0308), the Glozhene Formation (samples 0307–329), and the lower part of the Salash Formation (330–346). The sampling interval was usually 1 m, and was 5 m in the Salash Formation.

The two Gintsi sections (Text-figs 3, 4) are located in the southern part of the village of Gintsi, Sofia District, on the right bank of the Nishava River (Text-fig. 1). 50 thin sections, with a sampling interval of 1 m, were examined from the Gintsi 1 section, spanning the upper part of the Gintsi Formation (10 m), the Glozhene Formation (37 m), and the lowermost part of the Salash Formation (3 m) (Text-fig. 3). 54 thin sections from the Gintsi 2 section came from various sources; we used the old sample set of Bakalova (in Nikolov and Sapunov 1977) for the upper part of the Gintsi Formation (13 m, samples 608–610) and the lower part of the Glozhene Formation (19 m, samples 611–618). New samples were taken from the highest 7 m of the Glozhene Formation (samples 1–18), and from the basal 31 m of the lower part of the Salash Formation (samples 19–39) (Text-fig. 4). The average sample interval in the Gintsi 2 section was between 1 and 2 m.


The Gintsi 2 section is the type section of the Gintsi Formation (Sapunov and Ziegler 1976), and it is the section presented during the International symposium on the Jurassic/Cretaceous boundary in Bulgaria (Nikolov and Sapunov 1977).

SECTIONS STUDIED AND MATERIAL

The study is based on three sections, Barlya, Gintsi 1 and Gintsi 2, located in the Western Balkan Mts, western Bulgaria (Text-fig. 1).


CALPIONELLID BIOZONATION OF THE WESTERN BALKAN MTS

The microfossils of the succession studied are represented by calpionellids, calcareous nannofossils, cal-



Text-fig. 1. Location map of Bulgaria (courtesy of L. Metodiev) with position of the study area (A) and geological map (B) of the study area with positions of the sections studied

Gintsi 1 section



Text-fig. 3. Lithological log with calpionellid occurrences in the Gintsi 1 section


careous dinocysts, globochaetids, radiolarians and small benthic foraminifers. Macrofossils, such as bivalves, gastropods, ostracods, aptychi and juvenile ammonites, also occur. Spicules of siliceous sponges, crinoid and echinoid fragments derived from a neighbouring carbonate platform are less common. The fossil content identifies the sediments as hemipelagic to pelagic deposits. Koleva-Rekalova (in Lakova *et al.* 2007) classified the limestones as wackestones and

mudstones and distinguished the following microfacies in stratigraphic order: Saccocoma, Globochaete, Calpionellid and Spicule microfacies.

The calpionellids are characterized by rapid evolution and vast geographic distribution, so they are ideal for biostratigraphic studies. They are essential for fine subdivision, precise dating and reliable long-distance correlation of pelagic carbonates of Tithonian to Valanginian age throughout the Mediterranean Realm.

TITHONIAN TO VALNGINIAN CALPIONELLID ZONATION

Gintsi 2 section



Text-fig. 4. Lithological log with calpionellid occurrences in the Gintsi 2 section

Numerous calpionellid zonal schemes exist. The zonal and subzonal scheme of Pop (1994, 1997b) and Reháková and Michalík (1997a, b), slightly modified by Lakova (1993) and Lakova *et al.* (1999, 2007), is used herein. Seven calpionellid zones and eleven interval subzones are recognized and characterized in the present report (Text-fig. 6).


This zonation is based on the vertical ranges of 57 species of chitinoideids and calpionellids. Fourteen successive calpionellid bioevents, mainly first occurrences (FOs), serve as indicators of the lower zonal

and subzonal boundaries (Text-fig. 6). Thus, the calpionellid zones and subzones are almost invariably interval zones.

Tithonian

Chitinoidea Zone (Lower and Upper Tithonian); Plates 1, 5

The Chitinoidea Zone was introduced by Enay and Geysant (1975) and defined by Grandesso (1977).



Text-fig. 5. Calpionellid bioevents within the Crassicollaria and Calpionella zones

It is subdivided into the Dobeni and Boneti subzones, characterized by Grandesso (1977) and Borza (1984). The lower boundary of the zone is defined by the FO of microgranular-walled chitinoideids.

Correlations: The Chitinoideida Zone was recorded from Spain (Enay and Geysant 1975, Pruner *et al.* 2010), Slovakia (Borza 1984; Borza and Michalík 1986; Reháková 1995; Reháková and Michalík 1997a, b; Houša *et al.* 1999a, b; Michalík *et al.* 2009; Grabowski *et al.* 2010b), Hungary (Grabowski *et al.* 2010a), Romania (Pop 1986b, 1994, 1997b, 1998b), Poland (Pszczółkowski 1996; Pszczółkowski and Myczyński 2004; Grabowski and Pszczółkowski 2006), Ukraine (Reháková *et al.* 2011), Italy (Grandesso 1977; Channell and Grandesso 1987; Grün and Blau 1997; Andreini *et al.* 2007; Houša *et al.* 2004), Austria (Reháková *et al.* 1996, 2009; Lukeneder *et al.* 2010), Turkey (Altiner and Özkan 1991), Tunisia (Boughdiri *et al.* 2006, 2009; Sallouhi *et al.* 2011), Morocco (Benzaggagh and Atrops 1995; Benzaggagh *et al.* 2010), Cuba (Pszczółkowski 1999; Pszczółkowski and Myczyński 2010). The presence of the genus *Chitinoideida* was also reported from Germany (Doben 1963). In Bulgaria, the Chitinoideida Zone was recognized in the Western Balkan Mts (Bakalova 1977; Lakova 1993; Lakova *et al.* 1999), in the Western Fore-Balkan (Lakova *et al.* 1999), in the Central Fore-Balkan (Ivanova 1997) and from the subsurface of the Moesian Platform of northeast Bulgaria (Ivanova *et al.* 2002).

Sallouhi *et al.* (2011) proposed a new, more detailed subdivision of the classic Chitinoideida Zone. They introduced the Longicollaria Zone, equivalent of the Dobeni

subzone, and restricted their Chitinoideida Zone to the interval of the widely used Boneti Subzone. Our study has not confirmed the presence of *Dobeniella bermudezi* below the FO of *Chitinoideida boneti*. Further detailed studies outside North Africa may support Sallouhi's *et al.* (2011) chitinoideid subzonal division.

Dobeni Subzone. Its lower boundary coincides with the zonal base, which is indicated by the FO of *Longicollaria dobeni*. The chitinoideids are represented by *Borziella slovenica*, *Dobeniella colomi*, *Dobeniella tithonica*, *Daciella svinitensis*, and *Carpathella rumanica*. The subzone corresponds to the upper part of the Gintsi Formation. Its thickness is 5 to 8 m.

This subzone was previously reported from the Western Carpathians, Slovakia (Borza 1984; Borza and Michalík 1986; Reháková 1995, 2002; Reháková and Michalík 1997a, b; Houša *et al.* 1999a, b; Michalík *et al.* 2009; Grabowski *et al.* 2010b), Southern Carpathians, Romania (Pop 1994, 1997b, 1998b), Italy (Andreini *et al.* 2007), Spain (Pruner *et al.* 2010), Hungary (Grabowski *et al.* 2010a), Tunisia (Boughdiri *et al.* 2006, 2009), Morocco (Benzaggagh and Atrops 1995; Benzaggagh *et al.* 2010); Cuba (Pszczółkowski and Myczyński 2010); and Bulgaria (Western Balkan Mts and the Western Fore-Balkan; Lakova 1993; Lakova *et al.* 1999).

Boneti Subzone. The base of the subzone is defined by the FO of *Chitinoideida boneti*. The assemblage is dominated by *Chitinoideida boneti*; *Longicollaria insueta*,

Dobeniella cubensis, *Dobeniella bermudezi*, *Almajella cristobalensis* and *Daciella danubica* also occur. The subzone is 4.5 to 7 m thick and comprises the uppermost part of the Gintsi Formation and the base of the Glozhene Formation.

The Boneti Subzone was recorded from Slovakia (Borza 1984; Borza and Michalík 1986; Reháková 1995, 2002; Reháková and Michalík 1997a, b; Houša *et al.* 1999a, b; Michalík *et al.* 2009; Grabowski *et al.* 2010b), Poland (Grabowski and Pszczółkowski 2006), Romania (Pop 1997b, 1998b), Italy (Grün and Blau 1997; Andreini *et al.* 2007; Houša *et al.* 2004), Spain (Pruner *et al.* 2010), Hungary (Grabowski *et al.* 2010a), Austria (Reháková *et al.* 2009; Lukeneder *et al.* 2010), Tunisia (Boughdiri *et al.* 2006, 2009), Morocco (Benzaggagh and Atrops 1995; Benzaggagh *et al.* 2010); Cuba (Pszczółkowski and Myczyński 2010); and Bulgaria (Gintsi 2 section, Bakalova 1977; Western Balkan Mts., Lakova 1993; Central Fore-Balkan, Ivanova 1997; and Western Fore-Balkan, Lakova *et al.* 1999).

Praetintinnopsella Zone (Upper Tithonian); Plates 1, 5

The base of the zone is defined by the FO of *Praetintinnopsella andrusovi* and its top by the FO of representatives of the family Calpionellidae with hyaline calcite wall (Grandesso 1977). Its interval includes ca. 0.5 to 2.0 m of the lower part of the Glozhene Formation. *P. andrusovi* co-occurs with *Chitinoidea boneti*, *Chitinoidea hegarati*, and single atypical *Tintinnopsella carpathica*.

Correlations: The *Praetintinnopsella* Zone was recorded from Slovakia (Borza 1984; Borza and Michalík 1986; Reháková 1995; Reháková and Michalík 1997a; Michalík *et al.* 2009), Romania (Pop 1994, 1997b, 1998b), Poland (Pszczółkowski 1996; Grabowski and Pszczółkowski 2006), Hungary (Grabowski *et al.* 2010a), Ukraine (Reháková *et al.* 2011), Austria (Reháková *et al.* 2009; Lukeneder *et al.* 2010), and from Italy (Grandesso 1977; Andreini *et al.* 2007). It is also known as the “andrusovi subzone” of the *Chitinoidea* Zone in Italy (Grün and Blau 1997), Spain (Pruner *et al.* 2010), and Cuba (Pszczółkowski and Myczyński 2010). The genus *Praetintinnopsella* also occurs in the upper part of the *Chitinoidea* Zone in Turkey (Altiner and Özkan 1991), without defining a separate zone. In Bulgaria the *Praetintinnopsella* Zone was distinguished in the Barlya section in the Western Balkan Mts (Lakova 1993), in the Gintsi 1 and 2 sections (Lakova and Petrova 2009), in the Central Fore-Balkan (Ivanova 1997), and in the Western Fore-Balkan (Lakova *et al.* 1999).

Crassicollaria Zone (Upper Tithonian); Plates 1, 2, 5, 6

The zone was introduced as the “A Zone” by Remane (1963). The *Crassicollaria* Zone was defined by Allemann *et al.* (1971). Its lower boundary is defined by the FO of hyaline-walled calpionellids and its upper boundary by the “explosion” of the medium-sized spherical variety of *Calpionella alpina*. The zone is here divided into two subzones, Remanei and Massutiniana.

Correlations: The zone is widely known from the Tethyan Realm: in France (Remane 1963, 1971; Le Hégarat and Remane 1968; Cecca *et al.* 1989), Italy (Catalano and Liguori 1971; Channell and Grandesso 1987; Grün and Blau 1997; Andreini *et al.* 2007; Houša *et al.* 2004), Spain (Allemann *et al.* 1975; Pruner *et al.* 2010), Slovakia (Borza 1984; Borza and Michalík 1986; Reháková 1995; Reháková and Michalík 1997a, b; Houša *et al.* 1999a, b; Michalík *et al.* 2009; Grabowski *et al.* 2010b), Poland (Wierzbowski and Remane 1992; Pszczółkowski 1996; Pszczółkowski and Myczyński 2004; Grabowski and Pszczółkowski 2006), Romania (Pop 1974, 1986b, c, 1994, 1997b, 1998b; Barbu and Melinte-Dobrinescu 2008), Hungary (Grabowski *et al.* 2010a), Austria (Reháková *et al.* 1996, 2009; Lukeneder *et al.* 2010), Greece (Skourtsis-Coroneou and Solakius 1999), Turkey (Altiner and Özkan 1991), Iran (Azimi *et al.* 2008), Tunisia (Boughdiri *et al.* 2006, 2009). It is the same as the “A zone” of Remane (1963) in Tunisia (Ben Abdesselam-Mahdaoui *et al.* 2010), Morocco (Benzaggagh and Atrops 1995; Benzaggagh *et al.* 2010), Cuba (Pop 1976; Pszczółkowski *et al.* 2005; Pszczółkowski and Myczyński 2010) and Mexico (Trejo 1980; Adatte *et al.* 1994). In Bulgaria the *Crassicollaria* Zone was established in the Western Balkan Mts and the Western Fore-Balkan (Bakalova 1977; Lakova 1993; Lakova *et al.* 1999), in the Central Fore-Balkan (Ivanova 1997) and is known from the subsurface of the Moesian Platform in northeast Bulgaria (Ivanova *et al.* 2002). Lakova *et al.* (2009) reported the *Crassicollaria* Zone from eastern Serbia.

Remanei Subzone. The subzone was defined by Remane *et al.* (1986) as the lower subzone of the *Crassicollaria* Zone, equivalent to the A1 subzone of Remane (1963). Its lower boundary is marked by the almost simultaneous appearance of *Tintinnopsella remanei* and the typical small variety of *Tintinnopsella carpathica*. *Crassicollaria intermedia* first appears in this subzone. The subzone is 4 to 5 m thick in the Glozhene Formation in all three sections.

Correlations: The subzone is known from Romania (Pop 1986b, 1994), Slovakia (Reháková and Michalík

1997a, b; Houša *et al.* 1999a, b; Michalík *et al.* 2009; Grabowski *et al.* 2010b), Poland (Wierzbowski and Remane 1992; Pszczółkowski 1996; Grabowski and Pszczółkowski 2006), Italy (Grün and Blau 1997; Andreini *et al.* 2007; Houša *et al.* 2004), Hungary (Grabowski *et al.* 2010a), Austria (Reháková *et al.* 2009; Lukeneder *et al.* 2010), Spain (Pruner *et al.* 2010), Tunisia (Boughdiri *et al.* 2006, 2009; Sallouhi *et al.* 2011) and Cuba (Pszczółkowski and Myczyński 2010). In Bulgaria it is known from the Western Balkan Mts and Western Fore-Balkan (Lakova 1993; Lakova *et al.* 1999), as well as from the subsurface of the Moesian Platform (Ivanova *et al.* 2002). The Remanei Subzone was determined as the A1 Subzone in France (Remane 1963, 1971; Cecca *et al.* 1989), Turkey (Altiner and Özkan 1991) and Morocco (Benzaggagh and Atrops 1995; Benzaggagh *et al.* 2009). It is equivalent to the lower *Crassicollaria intermedia* Subzone sensu Pop (1974) in Romania, and sensu Skourtsis-Coroneou and Solakius (1999) in Greece.

Massutiniana Subzone. This subzone was introduced by Lakova (1993). It corresponds to the A2 + A3 subzones of Remane (1963) and to the *Intermedia* Subzone of Remane *et al.* (1986). The lower boundary of the *Massutiniana* Subzone is marked by the FOs of *Calpionella grandalpina* and *Calpionella alpina*. The subzone is characterized by high diversity of the genus *Crassicollaria* and a quantitative increase in calpionellids. *Crassicollaria massutiniana*, *C. brevis* and *C. parvula* first appear in this zone, and *Calpionella ellipticalpina* is limited to the zone (Text-fig. 5). The subzone is 5 to 6 m thick in the Glozhene Formation.

The *Massutiniana* Subzone was previously recorded in Bulgaria (Lakova 1993; Lakova *et al.* 1999; Ivanova *et al.* 2002). It is equivalent to the *Intermedia* + *Colomi* Subzones in Romania (Pop 1994), to the *Brevis* + *Colomi* Subzones in Slovakia (Reháková and Michalík 1997a, b; Michalík *et al.* 2009), to the A2 + A3 Subzones in France (Remane 1963, 1971; Cecca *et al.* 1989), Poland (Wierzbowski and Remane 1992), Turkey (Altiner and Özkan 1991) and Morocco (Benzaggagh and Atrops 1995; Benzaggagh *et al.* 2010), to the *Intermedia* + *Catalanoi* Subzones in Italy (Grün and Blau 1997) and to the *Intermedia* Subzone in Italy (Andreini *et al.* 2007; Houša *et al.* 2004), Poland (Pszczółkowski 1996; Grabowski and Pszczółkowski 2006), Slovakia (Houša *et al.* 1999a, b; Grabowski *et al.* 2010b), Hungary (Grabowski *et al.* 2010a), Austria (Reháková *et al.* 2009; Lukeneder *et al.* 2010), Spain (Pruner *et al.* 2010), Romania (Barbu and Melinte-Dobrinescu 2008), Tunisia (Boughdiri *et al.* 2006, 2009) and Cuba (Pszczółkowski and Myczyński

2010); to the *Brevis-Parvula* Subzone in Romania (Pop 1974); and to the *Intermedia* + *Brevis* Subzones in Greece (Skourtsis-Coroneou and Solakius 1999).

Berriasian

Calpionella Zone (Lower Berriasian); Plates 2, 6

The *Calpionella* Zone was introduced and defined by Allemann *et al.* (1971). Its lower boundary is marked by the “explosion” of *Calpionella alpina*, i.e. the medium-sized spherical variety of this species. This boundary was recommended as the base of the Berriasian Stage at the Lyon-Neuchâtel Colloquium in 1973 and the 32nd International Geological Congress in Florence, 2004. The zone was divided into three subzones (Remane *et al.* 1986): *Alpina*, *Remaniella* and *Elliptica*.

Correlations: The zone is known from Slovakia (Borza 1984; Borza and Michalík 1986; Reháková 1995, 2000a; Reháková and Michalík 1997a, b; Houša *et al.* 1999a, b; Michalík *et al.* 2009; Grabowski *et al.* 2010b), Poland (Pszczółkowski 1996; Pszczółkowski and Myczyński 2004; Boorova *et al.* 2004; Grabowski and Pszczółkowski 2006), Romania (Pop 1974, 1986b, c, 1994, 1997b, 1998b; Barbu and Melinte-Dobrinescu 2008), eastern Serbia (Lakova *et al.* 2009), Greece (Skourtsis-Coroneou and Solakius 1999), Italy (Channell and Grandesso 1987; Grün and Blau 1997; Houša *et al.* 2004; Andreini *et al.* 2007), Austria (Vašíček *et al.* 1999; Lukeneder and Reháková 2004, 2007; Reháková *et al.* 1996, 2009; Lukeneder *et al.* 2010), Hungary (Grabowski *et al.* 2010b), Spain (Pruner *et al.* 2010), Iran (Azimi *et al.* 2008), Tunisia (Boughdiri *et al.* 2006, 2009), Morocco (Benzaggagh and Atrops 1995; Benzaggagh *et al.* 2009), Cuba (Pop 1976; Pszczółkowski *et al.* 2005; Pszczółkowski and Myczyński 2010), Mexico (Adatte *et al.* 1994) and Bulgaria (Bakalova 1977; Lakova 1993; Lakova *et al.* 1999; Ivanova 1997; Ivanova *et al.* 2002). It is known as the B + C zones in France (Remane 1963, 1971; Le Hégarat and Remane 1968; Cecca *et al.* 1989), and Turkey (Altiner and Özkan 1991), as the B + C zones (*Calpionella alpina* + *Calpionella elliptica*) in Italy (Catalano and Liguori 1971), Spain (Allemann *et al.* 1975) and Poland (Wierzbowski and Remane 1992); and as the *Calpionella alpina* + *Calpionella elliptica* Subzones, in Mexico (Trejo 1980). Ben Abdesselam-Mahdaoui *et al.* (2011) recognized five successive subzones in the *Calpionella* Zone, namely B1, B2, B3, C1 and C2. Of these, their B2+B3 subzones equate by definition with the *Remaniella* Subzone; and their C2 Subzone with the *Elliptica* Subzone.

Alpina Subzone. This subzone was introduced and defined by Pop (1974). Its base is defined by the “explosion” of *Calpionella alpina*, and its top by the FO of the genus *Remaniella*. *Calpionella alpina* is the dominant species in this subzone. Three other bioevents within the Alpina Subzone could be used for a further subdivision. These are the acme of *Crassicollaria parvula*, and the FOs of *Calpionella minuta* and *Tintinnopsella doliformis*. Additional studies may confirm the usefulness and geographic record of these minor bioevents, which have been so far documented in Spain (Pruner *et al.* 2010). *Crassicollaria colomi* has been recorded below and above the lower boundary of the Alpina Subzone (Text-fig. 5) in this study, as well as in the Komshtitsa section earlier (Lakova 1993; Lakova *et al.* 2007). This record of *Cr. colomi* in the Alpina Subzone differs from data reported in the Western and Southern Carpathians and elsewhere in the Mediterranean Realm, where *Cr. colomi* was documented only in the highest part of the underlying Crassicollaria Zone and was even used to define the Colomi Subzone. The Alpina Subzone is normally 16 m thick (Glozhene Formation) in the sections studied.

Correlations: The subzone is known from Romania (Pop 1974, 1986b, c, 1994, 1997b, 1998b; Barbu and Melinte-Dobrinescu 2008), eastern Serbia (Lakova *et al.* 2009), Slovakia (Reháková 1995, 2000a; Reháková and Michalík 1997a, b; Houša *et al.* 1999a, b; Michalík *et al.* 2009; Grabowski *et al.* 2010b), Poland (Pszczółkowski 1996; Pszczółkowski and Myczyński 2004; Grabowski and Pszczółkowski 2006), Ukraine (Reháková *et al.* 2011), Italy (Grün and Blau 1997; Houša *et al.* 2004; Andreini *et al.* 2007), Hungary (Grabowski *et al.* 2010a), Austria (Vašíček *et al.* 1999; Reháková *et al.* 1996, 2009; Lukeneder *et al.* 2010), Tunisia (Boughdiri *et al.* 2006, 2009), Greece (Skourtsis-Coroneou and Solakius 1999), Iran (Azimi *et al.* 2008) and Cuba (Pop 1976; Pszczółkowski and Myczyński 2010). Its correlatives are: the lower *Calpionella alpina* Zone in Italy (Catalano and Liguori 1971; Allemann *et al.* 1975); the lower B Zone in France, Morocco, Turkey (Remane, 1963, 1971; Le Hégarat and Remane 1968; Cecca *et al.* 1989; Benzaggagh and Atrops 1995; Benzaggagh *et al.* 2010; Altiner and Özkan 1991), and the lower *Calpionella* Zone in Mexico (Trejo 1980). In Bulgaria it is known as the *Calpionella alpina* Subzone in the Western Balkan Mts, Western Fore-Balkan and from the subsurface of the Moesian Platform (Bakalova-Ivanova 1986; Lakova 1993; Lakova *et al.* 1999; Ivanova 1997; Ivanova *et al.* 2002).

Remaniella Subzone. This subzone was defined by Pop (1974), and its lower boundary was redefined by Remane *et al.* (1986). The base of the Remaniella Sub-

zone is defined by the FO of *Remaniella ferasini* and/or *R. duranddelgai* and its upper boundary by the FO of *Calpionella elliptica*. It is characterized by *R. ferasini*, *R. duranddelgai*, *Calpionella alpina*, *C. minuta*, *Crassicollaria parvula* and *Tintinnopsella carpathica* (small form). The abundance of calpionellids decreases rapidly in this subzone. It is 3 m thick in the Gintsi 1 and Barlya sections, 0.5 m thick in the Gintsi 2 section; and is limited to the Glozhene Formation.

Correlations: The subzone is known from Romania (Pop 1974, 1986b, c), Serbia (Lakova *et al.* 2009), Poland (Pszczółkowski 1996), Italy (Houša *et al.* 2004; Andreini *et al.* 2007), Morocco (Boughdiri *et al.* 2006, 2009), Greece (Skourtsis-Coroneou and Solakius 1999) and Cuba (Pop 1976). It is known as the Ferasini Subzone in Romania (Pop 1994, 1997b, 1998b; Barbu and Melinte-Dobrinescu 2008), Slovakia (Reháková 1995; Reháková and Michalík 1997a, b; Grabowski *et al.* 2010b), Poland (Pszczółkowski and Myczyński 2004; Grabowski and Pszczółkowski 2006), Ukraine (Reháková *et al.* 2011), Austria (Vašíček *et al.* 1999; Lukeneder and Reháková 2004; Reháková *et al.* 1996, 2009; Lukeneder *et al.* 2010). Its equivalents are: the upper B Zone in France (Remane 1963, 1971; Le Hégarat and Remane 1968) and Turkey (Altiner and Özkan 1991); the upper *Calpionella alpina* Zone in Italy (Catalano and Liguori 1971) and Spain (Allemann *et al.* 1975), as well as the upper *Calpionella alpina* Zone sensu Grün and Blau (1997). In Bulgaria it was reported by Bakalova-Ivanova (1986); Lakova (1993); Lakova *et al.* (1999); Ivanova (1997); Ivanova *et al.* (2002).

Elliptica Subzone. Catalano and Liguori (1971) introduced the zone as the “*Calpionella elliptica* Zone”. Pop (1974) redefined it, placing its base at the FO of *C. elliptica*, as the *Calpionella elliptica* Subzone, the latter being equivalent to Catalano and Liguori’s zone. In the present study the Elliptica Subzone has been used in the sense of Pop (1974). It is 2 to 4.5 m thick in the upper parts of the Glozhene Formation.

Correlations: The subzone is known from Romania (Pop 1974, 1986b, c), Serbia (Lakova *et al.* 2009), Slovakia (Reháková 1995; Reháková and Michalík 1997a, b), Poland (Wierzbowski and Remane 1992; Pszczółkowski 1996), Austria (Vašíček *et al.* 1999; Lukeneder and Reháková 2004, 2007; Reháková *et al.* 1996, 2009; Lukeneder *et al.* 2010), Morocco (Boughdiri *et al.* 2006, 2009), Greece (Skourtsis-Coroneou and Solakius 1999), Iran (Azimi *et al.* 2008), Cuba (Pop 1976; Pszczółkowski and Myczyński 2010) and Mexico (Trejo 1980). It is also known as the *Calpionella elliptica* Zone in Italy (Cata-

lano and Liguori 1971) and Spain (Allemann *et al.* 1975). Its correlatives are: the C Zone in France (Remane 1963, 1971; Le Hégarat and Remane 1968) and Turkey (Altiner and Özkan 1991); the Elliptica + Longa subzones in Romania (Pop 1994, 1997b; Barbu and Melinte-Dobrinescu 2008), the Elliptica + Cadischiana subzones in Italy (Grün and Blau 1997; Andreini *et al.* 2007), Poland (Pszczółkowski and Myczyński 2004; Grabowski and Pszczółkowski 2006), Slovakia (Grabowski *et al.* 2010b) and Hungary (Fözy *et al.* 2010). In Bulgaria it was reported from the Western Balkan Mts and the Western Fore-Balkan by Bakalova-Ivanova (1986), Lakova (1993) Lakova *et al.* (1999) and Petrova (2010); from the Central Fore-Balkan by Ivanova (1997); and from the Moesian Platform by Ivanova *et al.* (2002).

Calpionellopsis Zone (Upper Berriasian); Plates 3, 4, 7

The zone was introduced as the “D Zone” by Remane (in Le Hégarat and Remane 1968) and defined subsequently as the Calpionellopsis Standard Zone by Allemann *et al.* (1971). It is defined by the FO of *Calpionellopsis simplex* (base) and the FO of *Calpionellites darderi* (top). It is divided into three subzones, Simplex, Oblonga and Murgeanui.

Correlations: The zone is known from France (Le Hégarat and Remane 1968; Blanc *et al.* 1994), Spain (Allemann *et al.* 1975; Aguado *et al.* 2000), Italy (Catalano and Liguori 1971; Channell and Grandesso 1987; Grün and Blau 1997; Andreini *et al.* 2007), Slovakia (Borza 1984; Borza and Michalík 1986; Reháková 1995, 2000a; Reháková and Michalík 1997a, b), Poland (Wierzbowski and Remane 1992; Pszczółkowski 1996; Pszczółkowski and Myczyński 2004; Grabowski and Pszczółkowski 2006), Romania (Pop 1974, 1986b, 1994, 1997b; Barbu and Melinte-Dobrinescu 2008), Greece (Skourtsis-Coroneou and Solakius 1999), Austria (Vašíček *et al.* 1999; Lukeneder and Reháková 2004, 2007; Reháková *et al.* 1996), Hungary (Fözy *et al.* 2010), Tunisia (Ben Abdesselam-Mahdaoui *et al.* 2010), Turkey (Altiner and Özkan 1991), Cuba (Pop 1976; Pszczółkowski *et al.* 2005; Pszczółkowski and Myczyński 2010) and Mexico (Adate *et al.* 1994). It is equivalent to the Calpionellopsis simplex–Calpionellites darderi Zone of Azimi *et al.* (2008). In Bulgaria, the zone was reported by Bakalova-Ivanova (1986); Lakova (1993); Lakova *et al.* (1999); Ivanova (1997); Ivanova *et al.* (2002) and Petrova (2010).

Simplex Subzone. The lower boundary of the subzone is defined at the FO of *Calpionellopsis simplex* (Remane *et al.* 1986). It is equivalent to the D1 Subzone of Remane (1971).

Correlations: It is known as the Calpionellopsis simplex subzone in France (Le Hégarat and Remane 1968), Romania (Pop 1986b, c, 1994), Poland (Wierzbowski and Remane 1992; Pszczółkowski 1996; Pszczółkowski and Myczyński 2004; Grabowski and Pszczółkowski 2006), Slovakia (Reháková 1995, 2000a; Reháková and Michalík 1997a, b), Austria (Lukeneder and Reháková 2004, 2007), Italy (Grün and Blau 1997; Andreini *et al.* 2007), Greece (Skourtsis-Coroneou and Solakius 1999), Hungary (Fözy *et al.* 2010), Iran (Azimi *et al.* 2008), Cuba (Pop 1976; Pszczółkowski and Myczyński 2010) and Mexico (Adate *et al.* 1994). Its correlatives are: the lower Calpionellopsis simplex–Calpionellopsis oblonga Zone in Sicily, Italy (Catalano and Liguori 1971) and the lower Calpionellopsis simplex Subzone in Spain (Allemann *et al.* 1975), the D1 Subzone in Tunisia (Ben Abdesselam-Mahdaoui *et al.* 2011). In Bulgaria, the Simplex Subzone was reported in the West Balkan Mts and West Fore-Balkan by Lakova (1993); Lakova *et al.* (1999); Petrova (2010).

Oblonga Subzone. This subzone corresponds to the D₂ + D₃ subzones of Remane (Le Hégarat and Remane 1968). Remane *et al.* (1986) introduced it as the Oblonga Subzone. Pop (1986b) restricted the subzone to the interval between the FO of *Calpionellopsis oblonga* (base) and the FO of *Praecalpionellites murgeanui* (top), this restricted interval being used by the present authors herein. It is characterized by the highest diversity of calpionellid species throughout the entire time span of their existence. The FOs of *Calpionellopsis* sp. A, *Remaniella filipes* and *R. cadischiana*, and the abundance of *Lorenziella hungarica* and *L. plicata*, are documented in this subzone. The Oblonga Subzone is 3 to 15 m thick in the lowermost part of the Salash Formation.

Correlations: The Calpionellopsis oblonga Subzone sensu Pop (1986b, c) is known from Romania (Pop 1986b, c, 1994; Barbu and Melinte-Dobrinescu 2008), Slovakia (Reháková 1995; Reháková and Michalík 1997a, b), Greece (Skourtsis-Coroneou and Solakius 1999), Austria (Vašíček *et al.* 1999; Lukeneder and Reháková 2004, 2007), Hungary (Fözy *et al.* 2010), Iran (Azimi *et al.* 2008) and Cuba (Pszczółkowski and Myczyński 2010). It is known as the lower Calpionellopsis oblonga Subzone sensu Remane *et al.* (1986) in Poland (Pszczółkowski 1996; Pszczółkowski and Myczyński 2004; Grabowski and Pszczółkowski 2006), and Italy (Andreini *et al.* 2007). Its correlatives are: the D2 in Tunisia (Ben Abdesselam-Mahdaoui *et al.* 2011), the D2 + lower D3 in France (Le Hégarat and Remane 1968; Remane 1971), Turkey (Altiner and Özkan 1991), and Poland (Wierzbowski and Remane 1992); the middle

Calpionellopsis simplex–*Calpionellopsis oblonga* Zone in Italy (Catalano and Liguori 1971) and Spain (Alleman *et al.* 1975); the Oblonga + Filipescui Subzones in Italy (Grün and Blau 1997), and the middle *Calpionellopsis* Zone in Cuba (Pop 1976). In Bulgaria, the *Calpionellopsis* Zone was recorded from the Western Balkan Mts and the Western Fore-Balkan by Lakova *et al.* (1999, 2007); Ivanova *et al.* (2002); Petrova (2010).

Murgeanui Subzone. It was introduced and defined by Pop (1986b). Its base is defined by the FO of *Pracalpi-onellites murgeanui*. This subzone is documented only in the Gintsi 2 and Barlya sections. It is about 3 to 7 m thick, in the basal part of the Salash Formation.

Correlations: The subzone is known from Romania (Pop 1986b, c, 1994), Slovakia (Reháková 1995; Reháková and Michalík 1997a, b), Austria (Vašíček *et al.* 1999), Greece (Skourtsis-Coroneou and Solakius 1999), Cuba (Pszczółkowski and Myczyński 2010), and Bulgaria (Lakova *et al.* 1999, 2007).

Lower Valanginian

Calpionellites Zone; Plates 4, 7

Alleman *et al.* (1971) designated the E zone of Remane (1971) as the *Calpionellites* Standard Zone. Later Pop (1974) restricted the top of the zone to the LO of the genus *Calpionellites*, and this concept is used herein. The lower boundary is placed at the FO of *Calpionellites darderi*. The latter is also a primary criterion recommended for the definition of the base of the Valanginian Stage at the 32nd International Geological Congress in Florence, 2004. Pop (1994) subdivided the *Calpionellites* Zone into the Darderi and Major subzones.

Correlations: The zone is known from Slovakia (Borza 1984; Reháková 1995; Reháková and Michalík 1997a, b), Poland (Pszczółkowski and Myczyński 2004), Romania (Pop 1994), Italy (Channell and Grandesso 1987; Grün and Blau 1997; Andreini *et al.* 2007), Spain (Aguado *et al.* 2000), Austria (Vašíček *et al.* 1999; Lukeneder and Reháková 2004, 2007; Reháková *et al.* 1996), Hungary (Fözy *et al.* 2010), Cuba (Pszczółkowski and Myczyński 2010) and Mexico (Adatte *et al.* 1994). It is known as the E Zone in Turkey (Altiner and Özkan 1991) and Tunisia (Ben Abdesselam-Mahdaoui *et al.* 2010, 2011), as the *Calpionellites darderi* Subzone in Romania sensu Pop (Pop 1989) and in Mexico sensu Trejo (Trejo 1980). In Bulgaria Bakalova (1977) and Lakova *et al.* (1999, 2007) reported the zone from the Western Balkan Mts and the Western Fore-Balkan, and Ivanova

et al. (2002) from the subsurface of the Moesian Platform.

Darderi Subzone. The subzone was first mentioned by Trejo (1980), but formally introduced by Pop (1994). Its lower boundary is marked by the FO of *Calpionellites darderi* (Pop 1994). It is 6.5 to 13 m thick in the Salash Formation. *Calpionella minuta* and *Praecalpionellites murgeanui* disappear within this subzone.

Correlations: The subzone is known from Romania (Pop 1986a, 1994, 1997b), Slovakia (Reháková 1995; Reháková and Michalík 1997a, b), Poland (Pszczółkowski and Myczyński 2004), Italy (Grün and Blau 1997; Andreini *et al.* 2007), Spain (Aguado *et al.*, 2000), Austria (Reháková *et al.* 1996; Vašíček *et al.* 1999; Lukeneder and Reháková 2004, 2007), Hungary (Fözy *et al.* 2010) and Cuba (Pszczółkowski and Myczyński 2010). In Bulgaria, Lakova *et al.* (1999, 2007) and Petrova *et al.* (2010) reported the subzone from the Western Balkan Mts and the Western Fore-Balkan, and Ivanova *et al.* (2000, 2002) from the Western Srednogorie and the Moesian Platform.

Major Subzone. The subzone was established by Pop (1994). The lower boundary is defined by the FO of *Calpionellites major* and its upper boundary by the LO of the genus *Calpionellites*. The interval of the Major Subzone is about 18 m thick, in the Salash Formation. The LOs of *Praecalpionellites* sp. A, *Calpionellites coronatus* and *Calpionellites caravacaensis* are noted from this subzone.

Correlations: The subzone is known from Romania (Pop 1994, 1997b), Slovakia (Reháková 1995; Reháková and Michalík 1997a, b), Poland (Vašíček *et al.* 1999; Pszczółkowski and Myczyński 2004), Italy (Grün and Blau 1997; Andreini *et al.* 2007), Austria (Lukeneder and Reháková 2004, 2007) and Cuba (Pszczółkowski and Myczyński 2010). In Bulgaria, Lakova *et al.* (1999, 2007) and Petrova *et al.* (2010) reported the subzone from the Western Balkan Mts and the Western Fore-Balkan, and Ivanova *et al.* (2000, 2002) from the Western Srednogorie and the Moesian Platform.

Upper Valanginian and Hauterivian

Tintinnopsella Zone

The *Tintinnopsella* Zone was introduced and defined by Borza (1984). Its base is defined by the LO of *Calpionellites* and its top by the LO of the family Cal-

TITHONIAN TO VALNGINIAN CALPIONELLID ZONATION

tionellidae. The calpionellid association characterizing the zone is poor in species; *Tintinnopsella longa*, *Lorenziella hungarica*, *L. plicata* and *Remaniella cadischiana* occur in the lower part of the zone only, which is late Valanginian in age (Petrova *et al.* 2010). Upwards, only *T. carpathica* persisted in the Hauterivian part. Pop (1994) reported on the occurrence of *T. carpathica* in Hauterivian successions dated on ammonites and proposed to divide the Tintinnopsella Zone into the Cadischiana and Carpathica subzones.

Correlations: The zone is known from Romania (Pop 1989, 1994), Slovakia (Borza 1984; Reháková 1995; Reháková and Michalík 1997a,b), Poland (Pszczółkowski and Myczyński 2004), Italy (Grün and Blau 1997; Andreini *et al.* 2007), Spain (Granier *et al.* 1995), Austria (Reháková *et al.* 1996, Vašíček *et al.* 1999; Lukeneder and Reháková 2004, 2007; Reháková *et al.* 2006), Hungary (Fözy *et al.* 2010), Turkey (Altiner and Özkan 1991), and as the Tintinnopsella carpathica Zone in Mexico (Trejo 1980). In Bulgaria, Lakova *et al.* (1999, 2007) and Petrova *et al.* (2010) reported the subzone from the Western Balkan Mts and the Western Fore-Balkan, and Ivanova *et al.* (2000, 2002) from the Western Srednogie and the Moesian Platform.

DISCUSSION

Most of the calpionellid bioevents applied herein are widely used in calpionellid biostratigraphy (Remane 1971, 1986; Remane *et al.* 1986; Pop 1974, 1997b; Reháková 1995; Reháková and Michalík 1997a, b; Lakova *et al.* (1999); Andreini *et al.* 2007). The following calpionellid bioevents (mainly FOs), in ascending stratigraphic order, which are used for definition of the bases of calpionellid zones and subzones, are documented in the West Balkan Mts. (Text-fig. 6):

The FO of chitinoideids (*Longicollaria dobeni* and related species)

FO of *Chitinoidea boneti*

FO of *Praetintinnopsella andrusovi*

FO of hyaline-walled calpionellids (*Tintinnopsella carpathica* and *T. remanei*)

FO of *Calpionella grandalpina* and *Calpionella alpina*

The almost coeval explosion of *Calpionella alpina*, decline of the genus *Crassicollaria* and LO of *Calpionella elliptalpina*

FO of the genus *Remaniella* (*R. ferasini* and *R. duranddelgai*)

FO of *Calpionella elliptica* and almost coeval FO of large variety of *T. carpathica*

FO of *Calpionellopsis simplex*

FO of *Calpionellopsis oblonga*

FO of *Praecalpionellites murgeanui*

FO of *Calpionellites darderi*

FO of *Calpionellites major*

LO of the genus *Calpionellites*

Recent publications emphasize the need for detailed studies of calpionellid stratigraphical distribution and the relationships of calpionellids to other microfossils, such as calcareous nannoplankton, radiolarians and calcareous dinocysts. Research into the calpionellid biostratigraphy of the Upper Tithonian and Berriasian currently focuses on integrated bio- and magnetostratigraphy and the proposal of reliable criteria to define the base of the Berriasian (Channell and Grandesso 1987; Houša *et al.* 1999a, b, 2004; Michalík *et al.* 2009; Pruner *et al.* 2009; Grabowski and Pszczółkowski 2006; Grabowski *et al.* 2010a, b; Lukeneder *et al.* 2010; Fözy *et al.* 2010; Grabowski 2011; Wimbledon *et al.* 2011, Michalík and Reháková 2011).

There is a general agreement on the subzonal divisions of the Chitinoidea, Calpionella, Calpionellopsis and Calpionellites zones. Further study is needed of the calpionellid distribution in the Upper Tithonian in order to work out generally acceptable subzones of the Crassicollaria Zone (Text-fig. 5).

As long as the stage and substage boundaries of the Tithonian to Hauterivian in the Mediterranean Realm are based mainly on ammonites, the correlation between ammonite and calpionellid ranges is crucial in determining the chronostratigraphic value of the latter. A great amount of work has already been done (Le Hégarat and Remane 1968; Enay and Geysant 1975; Cecca *et al.* 1989; Wierzbowski and Remane 1992; Pop 1994; Tavera *et al.* 1994; Benzaggagh and Atrops 1995; Granier *et al.* 1995; Vašíček *et al.* 1999; Aguado *et al.* 2000; Skupien *et al.* 2003; Lukeneder and Reháková 2004, 2007; Boughdiri *et al.* 2006, 2009; Pruner *et al.* 2009; Benzaggagh *et al.* 2009; Petrova 2009, Petrova *et al.* 2011); however, a number of questions remain to be answered.

Chronostratigraphic importance of the FO of chitinoideids

The available data on the timing of the first appearances of chitinoideids (earliest *Longicollaria dobeni* and related species) are rather scarce. Pruner *et al.* (2009) reported what they claim to be the first chitinoideids from the upper Admirandum/Biruncinatum ammonite Zone; and Benzaggagh *et al.* (2010) from the upper part of the coeval Burkhardticerias Zone.

The Gintsi Formation in the Gintsi 2 section spans an

interval from the middle Kimmeridgian (*Crusolicerias divisum* ammonite Zone) to the upper Lower Tithonian (Sapunov in Nikolov and Sapunov 1977). We noted the first appearance of chitinoideids in sample 609 of the Gintsi Formation, which is 8 m above the top of the *Virgatosimoceras rothpletzi* ammonite Zone (Sapunov in Nikolov and Sapunov 1977). Olóriz and Tavera (1989) correlated the upper part of the *V. rothpletzi* Zone with the *Richteri* and *Admirandum/Biruncinatum* ammonite zones in Spain. The latter two ammonite zones are regarded as of late Early Tithonian age (Gradstein *et al.* 2004). The appearance of the first chitinoideids from the base of the *Admirandum/Biruncinatum* ammonite Zone (upper Ponti Zone in North Africa) was also reported by Reháková and Michalík (1997) and Grün and Blau (1997).

Base of the Upper Tithonian

Recent studies from Spain and Tunisia have demonstrated that the FO of *Chitinoideella boneti* coincides with the base of the *Simplisphintes* ammonite Subzone of the *Microcanthum* ammonite Zone (Pruner *et al.* 2009; Benzaggagh *et al.* 2010), the level which is regarded as the base of the Upper Tithonian (Gradstein *et al.* 2004). Similar dating of this calpionellid event was formerly reported from Morocco (Boughdiri *et al.* 2006). This dating is also confirmed by magnetostratigraphy (magnetostratigraphic M20n) and calcareous dinocysts (*Colomisphaera fortis*). The coincidence of parts of these events was demonstrated in the Gresten Klippenbelt in Austria (Reháková *et al.* 2009), the Western Balkan Mts in Bulgaria (Lakova *et al.* 1999), the Puerto Escaño section in Spain (Pruner *et al.* 2009), the Western Carpathians in Slovakia (Michalík *et al.* 2009) and in Morocco (Benzaggagh *et al.* 2010).

The *Crassicollaria* Zone and its subzones

Based on the FO of *Crassicollaria colomi*, in the uppermost part of the *Crassicollaria* Zone, Pop (1994) and Reháková and Michalík (1997) proposed the Colomi Subzone in the topmost part of the zone. In the Western Balkanides sections (Lakova, 1993; Lakova *et al.* 1999), *Crassicollaria colomi* was documented in the lower part of the *Calpionella* Zone (Alpina Subzone). These contradictory data could be due to particular facies features and/or a specific local longer range of *Cr. colomi* in the West Balkan Mts. area. The almost coeval FOs of *Calpionella grandalpina* and/or *Calpionella alpina* mark an important calpionellid bioevent in the Upper Tithonian that is used to define the base of the Massutiniana Subzone (Lakova 1993), which coincides with the Intermedia (= A2 + A3) Subzone of Remane (1986). Also ap-

pearing at that level are the nannofossil *Microstaurus chisti* and the calcareous dinocyst *Stomiosphaera proxima* (Lakova *et al.* 1999; Reháková 2000b). Importantly, the level coincides with the base of the M19r magnetic polarity chron (Houša *et al.* 1999a, b; Pruner *et al.* 2009). These biological and physical events should be considered in the context of multidisciplinary high-resolution stratigraphy of the Jurassic/Cretaceous boundary interval as possible criteria for the definition of the base of Berriasian.

The Tithonian-Berriasian boundary

The base of the Berriasian and, by definition, the base of the Cretaceous System, coincides in calpionellid terms with the base of the *Calpionella* Zone. The base of this zone is very close to the base of the *Berriasella jacobi* ammonite Zone/Subzone, which was provisionally recommended by the Second Cretaceous Symposium in Brussels in 1996 (Zakharov *et al.* 1996) and widely accepted by the Lower Cretaceous Ammonite Working Group (Reboulet and Klein 2009), as the base of the Cretaceous System. The combined ammonite and calpionellid biostratigraphy in the Tethyan Realm indicated that the explosion of the small variety of *Calpionella alpina* occurs at or just below the base of the *B. jacobi* Zone (Cecca *et al.* 1989; Aguado *et al.* 1994; Boughdiri *et al.* 2006; Benzaggagh *et al.* 2010). In terms of magnetostratigraphy, the base of the Berriasian is within the M19n chron (Grabowski 2011 and references therein).

Regional stratigraphic hiatus across the Early-Late Berriasian boundary

In the study area, a stratigraphic hiatus between the Glozhene and Salash Formations has been documented. It covers a time interval from the latest Early Berriasian to the early Late Berriasian. The presence of a stratigraphic hiatus between the Glozhene and Salash Formations was first recognised by Mandov (1971). That author accepted it as a boundary between the Jurassic and Cretaceous without palaeontological data. The calpionellid biostratigraphy suggests that the stratigraphic gap corresponds to the upper *Elliptica* Subzone, the whole *Simplex* Subzone and the lower *Oblonga* Subzone.

Base of the Valanginian

The Berriasian/Valanginian boundary is placed at the base of the *Calpionellites* Zone. Direct co-occurrence of calpionellids and calcareous dinocysts was reported from the Tithonian to Valanginian in Slovakia, Poland, Ukraine, Austria and Bulgaria (Reháková 2000a, b; Michalík *et al.*

TITHONIAN TO VALANGINIAN CALPIONELLID ZONATION

2009; Reháková and Wierzbowski 2005; Reháková *et al.* 2009, 2011; Lakova *et al.* 1999, 2007; Ivanova *et al.* 2006). The base of the Valanginian was defined at the FOs of the ammonite *Thurmanniceras pertransiens* and the calpionellid *Calpionellites darderi* (Aguado *et al.* 2000), a level which coincides with the FO of the calcareous dinocyst *Colomisphaera conferta* (Ivanova in Lakova *et al.* 2007; Ivanova in Petrova *et al.* 2011).

Unsolved problems in applying certain calpionellid bioevents to zonation include:

1. Tripartite or bipartite subdivision of the Crassicollaria Zone?
2. Problems with recognition of the Murgeanui Subzone, the latter being no longer important for definition of the base of the Valanginian.
3. Avoiding the use of species with controversial occurrences, such as *Dobeniella bermudezi*, *Crassicollaria colomi*, *Tintinnopsella longa*, *Lorenziella hungarica*, *Remaniella filipes cui* and *Remaniella cadischiana*, as indicators of subzonal bases.
4. Future direct correlations of the following calpionellid bioevents in the Upper Tithonian with ammonite, calcareous dinocysts, calcareous nannofossil bioevents and magnetic polarity chrons: the FO of *Chitinoïdella boneti*; the FO of *Tintinnopsella carpathica* and *Tintinnopsella remanei*; the FO of *Calpionella grandalpina*; the “explosion” of *Calpionella alpina*; and the FOs of *Calpionella minuta* and the genus *Remaniella*.

CONCLUSIONS

- 57 species belonging to 17 genera were recognized in the Tithonian–Valanginian succession of the three sections studied, Barlya, Gintsi 1 and Gintsi 2, located in the Western Balkan Mts, western Bulgaria. The family Chitinoïdellidae is represented by 17 species and 8 genera, and the family Calpionellidae by 41 species and 9 genera.
- Application of the updated taxonomy of the Chitinoïdellidae proposed by Pop (1997a, 1998a), Reháková (2002) and Sallouhi *et al.* (2011) resulted in the identification of the Tithonian genera *Almajella*, *Longicollaria*, *Borziella*, *Dobeniella*, *Daciella* and *Carpathella* for the first time in Bulgaria, as well as the species *Chitinoïdella hegarati*, known so far exclusively from Tunisia.
- 7 zones and 11 subzones (from upper Lower Tithonian to Upper Valanginian–Hauterivian) are defined, and their correlation to other areas is discussed. The chronostratigraphic position of the selected zones/subzones is discussed and corrected; the Boneti Subzone

and the Praetintinnopsella Zone are included in the Upper Tithonian.

- Two stage boundaries have been fixed in the Gintsi 1, Gintsi 2 and Barlya sections (Text-figs 2–4) – the Tithonian–Berriasian and Berriasian–Valanginian boundaries. The Tithonian–Berriasian boundary is drawn at the base of the *Calpionella* Zone at the “explosion” of *Calpionella alpina* and the LO of *Calpionella elliptalpina*; and the base of the Valanginian at the FO of *Calpionellites darderi*. In two of the sections (Gintsi 1 and 2), a stratigraphic gap between the Elliptica Subzone and the Oblonga Subzone (upper part) has been established.
- This review of the calpionellid zonation in numerous regions from Central America, northern Africa, Europe and the Middle East revealed the progress in detailed biostratigraphy and the necessity of a general consensus on a standard calpionellid bioevent/subzonal scale for the Tithonian to Valanginian time interval in the Tethyan Realm.

Acknowledgements

This study was undertaken at the Geological Institute, Sofia. The authors thank Lubomir Metodiev for permission to use his unpublished geological map. IL is grateful to Ivo Sapunov and Platon Tchoumatchenco (Geological Institute, Sofia) for the fruitful years of joint field work and stratigraphic discussions. The publication is funded by the Ministry of Science and Education, Sofia, contract № D02–802 (Project № BG 051PO001-3.3-05/0001 of the scheme “Science–Business”, Operative Programme “Development of Human Resources”). The authors are indebted to the journal reviewers, Daniela Reháková, Joachim Blau and Gloria Andreini, and to the journal editors for their critical comments and useful suggestions.

REFERENCES

- Aadate, T., Stinnesbeck, W. and Remane, J. 1994. The Jurassic/Cretaceous boundary in Northeastern Mexico. Confrontation and correlations by microfacies, clay mineral mineralogy, calpionellids and ammonites. *Géobios, Mémoire Spécial* **17**, 37–56.
- Aguado, R., Company, M. and Tavera, J.M. 2000. The Berriasian/Valanginian boundary in the Mediterranean region: new data from the Caravaca and Cehegín sections, SE Spain. *Cretaceous Research*, **21**, 1–21.
- Alleman, F., Catalano, R., Fares, F. and Remane, J. 1971. Standard calpionellid zonation (Upper Tithonian–Valanginian) of the Western Mediterranean province. In:

- Farinacci, A. (Ed.), Proceedings of the II Planktonic Conference, Roma 1970, 1337–1340.
- Allemann, F., Grün, W. and Wiedmann, L. 1975. The Berriasian of Caravaca (Prov. of Murcia) in the Subbetic zone of Spain and its importance for defining this stage and the Jurassic-Cretaceous boundary. *Mémoires du Bureau de Recherches Géologiques et Minières*, **86**, 14–22.
- Altiner, D. and Özkan, S. 1991. Calpionellid zonation in North-Western Anatolia (Turkey) and calibration of the stratigraphic ranges of some benthic foraminifers at the Jurassic-Cretaceous boundary. *Geologica Romana*, **27**, 215–235.
- Andreini, G., Caracuel, J.E. and Parisi, G. 2007. Calpionellid biostratigraphy of the Upper Tithonian-Upper Valanginian interval in Western Sicily (Italy). *Swiss Journal of Geosciences*, **100**, 179–198.
- Azimi, R., Seyed-Emami, K. and Sadeghi, A. 2008. Introducing of Calpionellid Zonation at the Jurassic-Cretaceous Boundary in the Shal Section (South-East of Khal-khal). *Geosciences Scientific Quarterly Journal*, **68**, 16–25.
- Bakalova, D. 1977. La succession à calpionelles de la coupe près du village de Ginci, Bulgarie du Nord-Ouest. *Comptes rendus de l'Académie bulgare des Sciences*, **30**, 423–426.
- Bakalova-Ivanova, D. 1986. Peculiarities of the *Calpionella* Zone in Bulgaria. *Acta Geologica Hungarica*, **29**, 89–92.
- Barbu, V. and Melinte-Dobrinescu, M.C. 2008. Latest Jurassic to earliest Cretaceous paleoenvironmental changes in the Southern Carpathians, Romania: regional record of the late Valanginian nitrification event. *Cretaceous Research*, **29**, 790–802.
- Ben Abdesselam-Mahdaoui, S., Benzaggagh, M., Bouhleb, S. and Razgallah, S. 2010. Nouvelle données biostratigraphiques sur les niveaux de passage Jurassique-Crétacé et les calcaires du Campanien dans le secteur de Hammam Zriba-Jebel Guebli (Tunisie septentrionale). *Geo-Eco-Trop*, **34**, 113–126.
- Ben Abdesselam-Mahdaoui, S., Benzaggagh, M., Razgallah, S., Rebah, A. and Rakia, B. 2011. Les associations des calpionelles du Berriasien et du Valanginien inférieur de la Tunisie septentrionale. Comparaison avec les associations du Rif externe (Maroc). *Comptes Rendus Palevol*, **10**, 527–535.
- Benzaggagh, M. and Atrpos, F. 1995. The Chitinoïdella and Crassicollaria zones (Tithonian) in the inner part of the Prerif (Morocco). New data and correlation with ammonite zones. *Comptes Rendus de l'Académie des Sciences Paris, Série II a*, **320**, 227–234.
- Benzaggagh, M., Cecca, F. and Rouquet, I. 2010. Biostratigraphic distribution of ammonites and calpionellids in the Tithonian of the internal Prerif (Msila area, Morocco). *Palaontologische Zeitschrift*, **84**, 301–315.
- Blanc, E., Bulot, L.G. and Paicheler, J.-C. 1994. La coupe de référence de Montbrun-les-Bains (Drôme, SE France): un stratotype potentiel pour la limite Berriasien-Valanginien. *Comptes Rendus de l'Académie des Sciences Paris, Série II*, **320**, 101–108.
- Boorová, D., Skupien, P. and Vašíček, Z. 2004. Biostratigraphy of the Lower Cretaceous limestones of the Godula facies of the Silesian Unit, Outer Western Carpathians. *Bulletin of Geosciences*, **79**, 121–131.
- Borza, K. 1984. The Upper Jurassic – Lower Cretaceous parabiostatigraphic scale on the basis of Tintinninae, Cadosinidae, Stomiosphaeridae, Calcisphaerulidae and other microfossils from the West Carpathians. *Geologický Zborník – Geologica Carpathica*, **35**, 539–550.
- Borza, K. and Michalík, J. 1986. Problems with delimitation of the Jurassic/Cretaceous boundary in the Western Carpathians. *Acta Geologica Hungarica*, **29**, 133–149.
- Boughdiri, M., Sallouhi, H., Maalaoui, K., Soussi, M. and Cordey, F. 2006. Calpionellid zonation of the Jurassic-Cretaceous transition in North-Atlasic Tunisia. Updated Upper Jurassic stratigraphy of the “Tunisian trough” and regional correlations. *Comptes Rendus Géosciences*, **338**, 1250–1259.
- Boughdiri, M., Sallouhi, H., Haddad, S., Cordey, F. and Soussi, M. 2009. Integrated biostratigraphy and regional correlations of Upper Jurassic-lowermost Cretaceous series in northern Tunisia. *GFF*, **131**, 71–81.
- Catalano, R. and Liguori, V. 1971. Facies a calpionelle della Sicilia Occidentale. In: Farinacci, A. (Ed.), Proceedings of the II Planktonic Conference, Roma 1970, 167–210.
- Cecca, F., Enay, R. and Le Hégarat, G. 1989. L'Ardésien (Titonique supérieur) de la région stratotypique: séries de référence et faunes (ammonites, calpionelles) de la bordure ardéchoise. *Documents Laboratoires de Géologie, Lyon*, **107**, 1–115.
- Channell, J.E.T. and Grandesso, P. 1987. A revised correlation of Mesozoic polarity chrons and calpionellid zones. *Earth and Planetary Science Letters*, **85**, 222–240.
- Doben, K. 1963. Über Calpionelliden an der Jura/Kreide-Grenze. *Mitteilungen der Bayerisches Staatssammlung für Paläontologie und historische Geologie*, **3**, 35–50.
- Enay, R. and Geysant, J.R. 1975. Faunes titoniques des chaînes bétiques (Espagne méridionale). *Mémoire B.R.G.M.*, **86**, 39–55.
- Fözy, I., Janssen, N.M.M., Price, G.D., Knauer, J. and Pálffy, J. 2010. Integrated isotope and biostratigraphy of a Lower Cretaceous section from the Bakony Mountains (Transdanubian Range, Hungary): A new Tethyan record of the Weissert event. *Cretaceous Research*, **31**, 525–545.
- Grabowski, J. 2011. Magnetostratigraphy of the Jurassic/Cretaceous boundary interval in the Western Tethys and its correlations with other regions: a review. *Volumina Jurassica*, **9**, 105–128.

TITHONIAN TO VALANGINIAN CALPIONELLID ZONATION

- Grabowski, J., Haas, J., Marton, E. and Pszczółkowski, A. 2010a. Magneto- and biostratigraphy of the Jurassic-Cretaceous boundary in the Lokut section (Transdanubian range, Hungary). *Studia Geophysica et Geodaetica*, **54**, 1–26.
- Grabowski, J., Michalík, J., Pszczółkowski, A. and Lintnerová, O. 2010b. Magneto- and isotope stratigraphy around the Jurassic/Cretaceous boundary in the Vysoká Unit (Malé Karpaty Mts, Slovakia): correlations and tectonic implications. *Geologica Carpathica*, **61**, 309–326.
- Grabowski, J. and Pszczółkowski, A. 2006. Magneto- and biostratigraphy of the Tithonian-Berriasian pelagic sediments of the Tatra Mountains (central Western Carpathians, Poland): sedimentary and rock magnetic changes at the Jurassic/Cretaceous boundary. *Cretaceous Research*, **27**, 398–417.
- Gradstein, F., Ogg, J., Smith, A., Bleeker, W. and Lourens, L. 2004. A new Geologic Time Scale with special reference to Precambrian and Neogene. *Episodes*, **27**, 83–100.
- Grandesso, P. 1977. Gli strati a Precalponellidi del Titoniano e i loro rapporti con il Rosso Ammonitico Veneto. *Memoire di Scienze Geologiche*, **32**, 1–15.
- Granier, B., Virgone, A., Busnardo, R. and Bulot, L.G. 1995. Des calponelles dans l'Hauterivien supérieur. Découverte exceptionnelle à Busot (Alicante, Espagne). *Comptes Rendus de l'Académie des Sciences Paris, Série II a*, **321**, 1179–1186.
- Grün, B. and Blau, J. 1997. New aspects of calponellid biochronology: proposal for a revised calponellid zonal and subzonal division. *Revue de Paléobiologie*, **16**, 197–214.
- Houša, V., Krs, M., Krsova, M., Man, O., Pruner, P. and Vendohová, D. 1999a. High-resolution magnetostratigraphy and micropalaeontology across the J/K boundary strata at Brodno near Zilina, western Slovakia: Summary of results. *Cretaceous Research*, **20**, 699–717.
- Houša, V., Krs, M., Man, O., Pruner, P. and Vendohová, D. 1999b. Correlation of magnetostratigraphy and calponellid biostratigraphy of the Jurassic/Cretaceous boundary strata in the Western Carpathians. *Geologica Carpathica*, **50**, 125–144.
- Houša, V., Krs, M., Man, O., Pruner, P., Vendohová, D., Cecca, F., Nardi, G. and Piscitello, M. 2004. Combined magnetostratigraphic, palaeomagnetic and calponellid investigations across Jurassic/Cretaceous boundary strata in the Bosso Valley, Umbria, central Italy. *Cretaceous Research*, **25**, 771–785.
- Ivanova, D. 1997. Upper Jurassic zonation on cadosinids, stomiosphaerids and calponellids of the Central Forebalkan, Bulgaria. *Geologica Balcanica*, **27**, 33–47.
- Ivanova, D., Koleva-Rekalova, E., Lakova, I. and Metodiev, L. 2006. Callovian to Berriasian pelagic carbonates in the Western Fore-Balkan, Bulgaria: microfacies, ammonite and microfossil zonations. *Volumina Jurassica*, **4**, 175–177.
- Ivanova, D., Lakova, I., Polishina, P. and Koleva-Rekalova, E. 2002. Joint biostratigraphy and lithofacies of Berriasian and Valanginian limestones from subsurface sections in NE Bulgaria. *Geologica Balcanica*, **32**, 63–67.
- Ivanova, D., Stoykova, K. and Lakova, I. 2000. New microfossil data on the age relationship between Slivnitsa and Salash Formation in Dragoman region, Western Bulgaria. *Comptes rendus de l'Académie bulgare des Sciences*, **53**, 77–81.
- Lakova, I. 1993. Middle Tithonian to Berriasian praecalponellid and calponellid zonation of the Western Balkanides, Bulgaria. *Geologica Balcanica*, **23**, 3–24.
- Lakova, I. 1994. Numerical criteria of precise delimitation of the calponellid *Crassicollaria* and *Calponella* Zones in relation to the Jurassic/Cretaceous system boundary. *Geologica Balcanica*, **24**, 23–30.
- Lakova, I., Rabrenović, D. and Petrova, S. 2009. Calponellid biostratigraphy across the Jurassic-Cretaceous boundary in the Western Balkan Mountains (Bulgaria and Serbia). In: The 8th symposium of IGCP 506 “Marine and non-marine Jurassic: global correlation and major geological events”, Bucharest, 28 August-3 September 2009, Abstracts and field guide, 18–19.
- Lakova, I., Stoykova, K. and Ivanova, D. 1999. Calponellid, nannofossil and calcareous dinocyst bioevents and integrated biochronology of the Tithonian to Valanginian in the Western Balkanides, Bulgaria. *Geologica Carpathica*, **50**, 131–168.
- Lakova, I., Tchoumatchenco, P., Ivanova, D. and Koleva-Rekalova, E. 2007. Callovian to Lower Cretaceous pelagic carbonates in the West Balkan Mountain (Komshtitsa and Barlya sections): integrated biostratigraphy and microfacies. *Geologica Balcanica*, **36**, 81–89.
- Le Hégarat, G. and Remane, J. 1968. Tithonique supérieur et Berriasien de la bordure cevenole. Correlation des Ammonites et des Calponnelles. *Geobios*, **1**, 7–70.
- Lukeneder, A., Halássová, E., Kroh, A., Mayrhofer, S., Pruner, P., Reháková, D., Schnabl, P., Sprovieri, M. and Wargreich, M. 2010. High resolution stratigraphy of the Jurassic-Cretaceous boundary interval in the Gresten Klippenbelt (Austria). *Geologica Carpathica*, **61**, 365–381.
- Lukeneder, A. and Reháková, D. 2004. Lower Cretaceous section of the Ternberg Nappe (Northern calcareous Alps, Upper Austria): facies-changes, biostratigraphy and paleoecology. *Geologica Carpathica*, **55**, 227–237.
- Lukeneder, A. and Reháková, D. 2007. Chronostratigraphic significance of an early Valanginian (Cretaceous) calponellid association (Hochkogel section, Upper Austria, Northern Calcareous Alps). *Geological Quarterly*, **51**, 27–38.

- Mandov, G. 1971. New stratigraphic data on the Lower Cretaceous in the Gubesh Syncline. *Annual of Sofia University, Faculty of Geology and Geography*, **63**, 47–60. [In Bulgarian with French abstract]
- Michalík, J. and Reháková, D. 2011. Possible markers of the Jurassic/Cretaceous boundary in the Mediterranean Tethys: A review and state of art. *Geoscience Frontiers*, **2**, 475–490.
- Michalík, J., Reháková, D., Halásová, E. and Lintnerová, O. 2009. The Brodno section – a potential regional stratotype of the Jurassic/Cretaceous boundary (Western Carpathians). *Geologica Carpathica*, **60**, 213–232.
- Nikolov, T. and Sapunov, I. 1970. On the regional stratigraphy of Upper Jurassic and Lower Cretaceous in Balkanides. *Comptes rendus de l'Académie bulgare des Sciences*, **23**, 1397–1400. [In Russian]
- Nikolov, T. and Sapunov, I. 1977. Excursion Guidebook. International symposium on the Jurassic/Cretaceous boundary in Bulgaria, 1–127 pp. Sofia University Press; Sofia.
- Nikolov, T. and Tsankov, T. 1971. Lithostratigraphic notes on part of the Lower Cretaceous sediments in the Western Balkanides. *Proceedings of the Geological Institute, Bulgarian Academy of Sciences, Series Stratigraphy and Lithology*, **20**, 63–70. [In Bulgarian]
- Olóriz, F. and Tavera, J.M. 1989. The significance of Mediterranean ammonites with regard to the traditional Jurassic–Cretaceous boundary. *Cretaceous Research*, **10**, 221–237.
- Petrova, S. 2009. Valanginian ammonite biostratigraphy in the Barlya section, Salash Formation (Western Balkan Mountains). Proceedings of the National Conference “Geosciences – 2009”, Sofia, 3–4 December, 59–60.
- Petrova, S. 2010. Calpionellid data on Berriasian age of the Sarbenitsa Wedge, Cherni Osam Formation (Western Balkan Mountains, Bulgaria). *Comptes rendus de l'Académie bulgare des Sciences*, **63**, 733–740.
- Petrova, S., Lakova, I. and Ivanova, D. 2010. The Salash Formation in Western Srednogorie: lithology, age and boundaries. Proceedings of the National Conference “Geosciences – 2010”, Sofia, 9–10 December, 93–94. [In Bulgarian]
- Petrova, S., Lakova, I. and Ivanova, D. 2011. Berriasian – Valanginian boundary in Bulgaria, *Review of the Bulgarian Geological Society*, **72**, 91–97.
- Pop, G. 1974. Les zones des Calpionelles Tithonique-Valanginiens du silon de Reșița (Carpathes meridionales). *Revue Roumaine de Géologie Géophysique et Géographie, Série Géologie*, **18**, 109–125.
- Pop, G. 1976. Tithonian – Valanginian calpionellid zones from Cuba. *Dări de Seamă ale Ședințelor Institutul de Geologie și Geofizică, 3. Paleontologie*, **62**, 237–266.
- Pop, G. 1986a. Réflexions sur certaines calpionelles Néocomiennes. *Dări de Seamă ale Ședințelor, Institutul de Geologie și Geofizică, 3. Paleontologie*, **70-71**, 103–108.
- Pop, G. 1986b. Les zones des calpionelles Tithonique-Néocomiennes de la région de Svinîța (Carpathes méridionales). *Dări de Seamă ale Ședințelor, Institutul de Geologie și Geofizică. 4. Stratigraphie*, **70-71**, 87–108.
- Pop, G. 1986c. Calpionellids and correlation of Tithonian-Valanginian formations. *Acta Geologica Hungarica*, **29**, 93–102.
- Pop, G. 1989. Age and facies of the calpionellid formations from the South Carpathians. In: Wiedmann, J. (Ed.), Cretaceous of the Western Tethys. Proceedings 3rd International Cretaceous Symposium, Tübingen 1987, 525–542, Schweizerbart Verlag, Stuttgart.
- Pop, G. 1994. Calpionellid evolutive events and their use in biostratigraphy. *Romanian Journal of Stratigraphy*, **76**, 7–24.
- Pop, G. 1997a. Révision systématique des chitinoïdelles Tithoniennes des Carpathes méridionales (Roumanie). *Comptes Rendus de l'Académie des Sciences Paris, Série II a*, **324**, 931–938.
- Pop, G. 1997b. Tithonian to Hauterivian praecalpionellids and calpionellids: bioevents and biozones. *Mineralia Slovaca*, **29**, 304–305.
- Pop, G. 1998a. Nouvelles chitinoïdelles Tithoniennes des Carpathes méridionales (Roumanie). *Comptes Rendus de l'Académie des Sciences, Paris, Série de la terre et des planètes*, **326**, 817–822.
- Pop, G. 1998b. Stratigraphic distribution and biozonation of Tithonian praecalpionellids and calpionellids from the South Carpathians. *Romaian Journal of Stratigraphy*, **77**, 3–25.
- Pszczółkowski, A. 1996. Calpionellid stratigraphy of the Tithonian-Berriasian pelagic limestones in the Tatra Mts (Western Carpathians). *Studia Geologica Polonica*, **109**, 103–130.
- Pszczółkowski, A. 1999. New data on the Lower Cretaceous microfossil and nannoconid stratigraphy in the Guaniguanico terrane of western Cuba. *Studia Geologica Polonica*, **114**, 7–33.
- Pszczółkowski, A., Garcia Delgado, D. and Gil Gonzalez, S. 2005. Calpionellid and nannoconid stratigraphy and microfacies of limestones at the Tithonian-Berriasian boundary in the Sierra del Infierno (Western Cuba). *Anales Societatis Geologorum Poloniae*, **75**, 1–16.
- Pszczółkowski, A. and Myczyński, R. 2004. Ammonite-supported microfossil and nannoconid stratigraphy of the Tithonian-Hauterivian limestones in selected sections of the Branisko Succession, Pieniny Klippen Belt (Poland). *Studia Geologica Polonica*, **123**, 133–197.
- Pszczółkowski, A. and Myczyński, R. 2010. Tithonian-Early

TITHONIAN TO VALNGINIAN CALPIONELLID ZONATION

- Valanginian evolution of deposition along the proto-Caribbean margin of North America recorded in Guaniguanico successions (western Cuba). *Journal of South American Earth Sciences*, **29**, 225–253.
- Pruner, P., Houša, V., Olóriz, F., Košťák, M., Krs, M., Man, O., Schanbl, P., Venhodová, D., Tavera, J.M. and Mazuch, M. 2010. High-resolution magnetostratigraphy and biostratigraphic zonation of the Jurassic-Cretaceous boundary strata in the Puerto Escano section (southern Spain). *Cretaceous Research*, **31**, 192–206.
- Reboullet, S., Klein, J. (reporters), Barragán, R., Company, M., González-Arreola, C., Lukeneder, A., Raisosadat, S., Sandoval, J., Szives, O., Tavera, J., Vašíček, Z. and Vermeulen, J. 2009. Report on the 3rd International Meeting of the IUGS Lower Cretaceous Ammonite Working Group, the “Kilian Group” (Vienna, Austria, 15th April 2008). *Cretaceous Research*, **30**, 496–502.
- Reháková, D. 1995. Nové poznatky o distribúci kalpionelíd vo vrchnojurských a spodnokriedových súvrstviach Západných Karpát. *Mineralia Slovaca*, **27**, 308–318.
- Reháková, D. 2000a. Calcareous dinoflagellate and calpionellid bioevents versus sea-level fluctuations recorded in the West-Carpathian (Late Jurassic/Early Cretaceous) pelagic environments. *Geologica Carpathica*, **51**, 229–243.
- Reháková, D. 2000b. Evolution and distribution of the Late Jurassic and Early Cretaceous calcareous dinoflagellates recorded in the Western Carpathian pelagic carbonate facies. *Mineralia Slovaca*, **32**, 79–88.
- Reháková, D. 2002. *Chitinoïdella* Trejo, 1975 in middle Tithonian Carbonate pelagic sequences of the West Carpathian Tethyan area. *Geologica Carpathica*, **53**, 369–379.
- Reháková, D., Halásová, E. and Lukeneder, A. 2009. The Jurassic-Cretaceous boundary in the Gresten Klippenbelt (Nutzhof, Lower Austria): Implications for Micro- and Nanofacies analysis. *Annalen des Naturhistorischen Museums in Wien*, **110 A**, 345–381.
- Reháková, D., Matyja, B., Wierzbowski, A., Schlögl, J., Krobicki, M. and Barski, M. 2011. Stratigraphy and microfacies of the Jurassic and lowermost Cretaceous of the Veliky Kamenets section (Pieniny Klippen Belt, Carpathians, Western Ukraine). *Volumina Jurassica*, **9**, 61–104.
- Reháková, D. and Michalík, J. 1997a. Calpionellid associations versus late Jurassic and Early Cretaceous sea – level fluctuations. *Mineralia Slovaca*, **29**, 306–307.
- Reháková, D. and Michalík, J. 1997b. Evolution and distribution of calpionellids – the most characteristic constituents of Lower Cretaceous Tethyan microplankton. *Cretaceous Research*, **18**, 495–504.
- Reháková, D., Michalík, J. and Ožvoldová, L. 1996. New microbiostratigraphical data from several Lower Cretaceous pelagic sequences of the Northern Calcareous Alps, Austria (preliminary results). *Geologisch-Paläontologische Mitteilungen Innsbruck*, **4**, 57–81.
- Reháková, D. and Wierzbowski, A. 2005. Microfacies and stratigraphic position of the Upper Jurassic Rogoża coquinas at Rogoźnik, Pieniny Klippen Belt, Carpathians. *Volumina Jurassica*, **3**, 15–27.
- Remane, J. 1963. Les Calpionelles dans les couches de passage jurassique-crétacé de la fosse vocontienne. *Travaux du Laboratoire de Géologie de la Faculté des Sciences de Grenoble*, **39**, 25–82.
- Remane, J. 1971. Les Calpionelles, Protozoaires planctoniques des mers mésogéennes de l'époque secondaire. *Annales Guébhard*, **47**, 369–432.
- Remane, J. 1986. Calpionellids and the Jurassic-Cretaceous boundary. *Acta Geologica Hungarica*, **29**, 15–26.
- Remane, J., Bakalova-Ivanova, D., Borza, K., Knauer, J., Nagy, I., Pop, G. and Tardi-Filacz, E. 1986. Agreement on the subdivision of the standard calpionellid zones defined at the II planktonic conference, Roma 1970. *Acta Geologica Hungarica*, **29**, 5–14.
- Sapunov, I. 1976. Ammonite stratigraphy of the Upper Jurassic in Bulgaria. I. Rock and ammonite successions. *Geologica Balcanica*, **6**, 17–40.
- Sapunov, I. and Ziegler, B. 1976. Stratigraphische Probleme im Oberjura des westlichen Balkangebirges. *Stuttgarter Beiträge zur Naturkunde, Serie B*, **118**, 1–46.
- Sallouhi, H., Boughdiri, M. and Cordey, F. 2011. Tithonian Chitinoïdellids of the South-Tethyan Margin of the Maghreb: New data from Tunisia. *Comptes Rendus Palevol*, **10**, 641–653.
- Skourtsis-Coroneou, V. and Solakius, N. 1999. Calpionellid zonation at the Jurassic/Cretaceous boundary within the Vigla Limestone Formation (Ionian Zone, western Greece) and carbon isotope analysis. *Cretaceous Research*, **20**, 583–595.
- Skupien, P., Vašíček, Z., Reháková, D. and Halásová, E. 2003. Biostratigraphy of Lower Cretaceous of the Manín unit (Butkov quarry, Strážovské Vrchy Mts., Western Carpathians). *Sborník vědeckých prací Vysoké školy báňské – Technické univerzity Ostrava, Řada hornicko-geologická*, **49**, 91–98.
- Tavera, J.M., Aguado, R., Company, M. and Olóriz, F. 1994. Intergated biostratigraphy of the Durangites and Jacobi Zones (J/K boundary) at the Puerto Escaño section in southern Spain (Province of Cordoba). *Geobios, Mémoire Special*, **17**, 469–476.
- Trejo, M. 1980. Distribucion estratigrafica de los Tintinidos mesozoicos mexicanos. *Revista del Instituto Mexicano del Petroleo*, **12**, 4–13.
- Vašíček, Z., Reháková, D. and Faupl, P. 1999. Zur Biostratigraphie der Schrambachschichten der Oisbergmulde bei Hollenstein a. d. Ybbs (Lunzer Decke, Kalkalpen, Nie-

- derösterreich). *Abhandlungen der Geologischen Bundesanstalt-A.*, **56**, 625–650.
- Wierzbowski, A. and Remane, J. 1992. The ammonite and calpionellid stratigraphy of the Berriasian and lowermost Valanginian in the Pieniny Klippen Belt (Carpathians, Poland). *Eclogae Geologicae Helvetiae*, **85**, 871–891.
- Wimbledon, W.A.P., Casselato, C.E., Reháková, D., Bulot, L.G., Erba, E., Gardin, S., Verreussel, R.M.C.H., Munstermann, D.K. and Hunt, C.O. 2011. Fixing a basal Berriasian and the Jurassic – Cretaceous (J-K) boundary – is there perhaps some light at the end of the tunnel? *Rivista Italiana di Paleontologia e Stratigrafia*, **117**, 295–307.
- Zakharov, V.A., Bown, P. and Rawson, P.F. 1996. The Berriasian Stage and the Jurassic-Cretaceous boundary. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Sciences de la Terre, SUPPL.*, **66**, 7–10.

Manuscript submitted: 27th August 2012

Revised version accepted: 15th March 2013

TITHONIAN TO VALNGINIAN CALPIONELLID ZONATION

APPENDIX

List of chitinoideid and calpionellid species

- Almajella cristobalensis* (Furrazola-Bermúdez, 1965)
Borziella slovenica (Borza, 1969)
Carpathella rumanica Pop, 1998
Chitinoideidella boneti Doben, 1963
Chitinoideidella elongata Pop, 1997
Chitinoideidella hegarati Sallouhi, Boughdiri and Cordey, 2011
Daciella danubica Pop, 1998
Daciella svinitensis Pop, 1998
Daciella sp. A
Dobeniella bermudezi (Furrazola-Bermúdez, 1965)
Dobeniella colomi (Borza, 1966)
Dobeniella cubensis (Furrazola-Bermúdez, 1965)
Dobeniella cf. cubensis (Furrazola-Bermúdez, 1965)
Dobeniella tithonica (Borza, 1969)
Longicollaria dobeni (Borza, 1966)
Longicollaria insueta (Řehánek, 1986)
Praetintinnopsella andrusovi Borza, 1969
Calpionella alpina Lorenz, 1902
Calpionella elliptalpina Nagy, 1986
Calpionella elliptica Cadisch, 1932
Calpionella grandalpina Nagy, 1986
Calpionella minuta Houša, 1990
Calpionella sp. A
Calpionellites caravacaensis Allemann, 1975
Calpionellites coronatus Trejo, 1975
Calpionellites darderi (Colom, 1934)
Calpionellites major (Colom, 1948)
Calpionellopsis oblonga (Cadisch, 1932)
Calpionellopsis simplex (Colom, 1939)
Calpionellopsis sp. A
Crassicollaria brevis Remane, 1962
Crassicollaria colomi Doben, 1963
Crassicollaria intermedia (Durand-Delga, 1957)
Crassicollaria massutiniana (Colom, 1948)
Crassicollaria parvula Remane, 1962
Lorenziella hungarica Knauer and Nagy, 1964
Lorenziella plicata Remane, 1968
Praecalpionellites hillebrandti Grün and Blau, 1999
Praecalpionellites murgeanui (Pop, 1974)
Praecalpionellites siriniaensis Pop, 1986
Praecalpionellites sp. A
Remaniella borzai Pop, 1994
Remaniella cadischiana (Colom, 1948)
Remaniella catalanoi Pop, 1996
Remaniella colomi Pop, 1996
Remaniella duranddelgai Pop, 1996
Remaniella ferasini (Catalano, 1965)
Remaniella filipescui Pop, 1994
Sturiella oblonga Borza, 1981
Sturiella aff. *oblonga* Borza, 1981
Tintinnopsella carpathica (Murgeanu and Filipescu, 1933)
Tintinnopsella aff. *carpathica* (Murgeanu and Filipescu, 1933)
Tintinnopsella dacica Filipescu and Dragastan, 1970
Tintinnopsella doliphormis (Colom, 1939)
Tintinnopsella longa (Colom, 1939)
Tintinnopsella remanei Borza, 1969
Tintinnopsella subacuta (Colom, 1948)

PLATE 1

Calpionellids of the Barlya section, Western Balkan Mts.

1-5 – *Daciella danubica* Pop, Lower Tithonian, Chitinoidea Zone, Dobeni Subzone, sample 0310. **6-8** – *Daciella svinitensis* Pop, Lower Tithonian, Chitinoidea Zone, Dobeni Subzone, **6** – sample 0310, **7-8** – sample 0309. **9-10** – *Borziella slovenica* (Borza), Lower Tithonian, Chitinoidea Zone, Dobeni Subzone, sample 0309. **11-12** – *Longicollaria dobeni* (Borza), Lower Tithonian, Chitinoidea Zone, Dobeni Subzone, sample 0309. **13-16** – *Dobeniella colomi* (Borza), Lower Tithonian, Chitinoidea Zone, Dobeni Subzone, **13** – sample 0310, **14-16** – sample 0309. **17-18** – *Chitinoidea boneti* Döben, Upper Tithonian, Chitinoidea Zone, Boneti Subzone, sample 0303. **19** – *Chitinoidea hegarati* Sallouhi, Boughdiri and Cordey, Upper Tithonian, Praetintinnopsella Zone, sample 0301. **20-21** – *Chitinoidea elongata* Pop, Upper Tithonian, Chitinoidea Zone, Boneti Subzone, **20** – sample 0302, **21** – sample 0303. **22**. *Almajella cristobalensis* (Furrázola-Bermúdez), Upper Tithonian, Chitinoidea Zone, Boneti Subzone, sample 0303. **23-24** – *Dobeniella cf. cubensis* (Furrázola-Bermúdez), Upper Tithonian, Chitinoidea Zone, Boneti Subzone, sample 0301. **25-26** – *Dobeniella bermudezi* (Furrázola-Bermúdez), Upper Tithonian, Chitinoidea Zone, Boneti Subzone, **25** – sample 0302; Praetintinnopsella Zone, **26** – sample 300. **27-31** – *Praetintinnopsella andrusovi* Borza, Upper Tithonian, Praetintinnopsella Zone, **27-29, 31** – sample 0301, **30** – sample 300. **32-33** – *Crassicollaria intermedia* (Durand-Delga), Upper Tithonian, Crassicollaria Zone, Remanei Subzone, **32** – sample 301, **33** – sample 302. **34** – *Tintinnopsella remanei* Borza, Upper Tithonian, Crassicollaria Zone, Remanei Subzone, sample 301. **35-36** – *Tintinnopsella carpathica* (Murgeanu & Filipescu), Upper Tithonian, Praetintinnopsella Zone, sample 300.




PLATE 2

Calpionellids of the Barlya section, Western Balkan Mts.

1 – *Crassicollaria massutiniana* (Colom), Upper Tithonian, Crassicollaria Zone, Massutiniana Subzone, sample 304-4. **2-3** – *Crassicollaria brevis* Remane, Upper Tithonian, Crassicollaria Zone, Massutiniana Subzone, sample 307. **4-5** – *Crassicollaria parvula* Remane, Upper Tithonian, Crassicollaria Zone, Massutiniana Subzone, **4** – sample 307, **5** – sample 310. **6-7** – *Crassicollaria colomi* Doben, Lower Berriasian, Calpionella Zone, Alpina Subzone, sample 313. **8-9** – *Calpionella grandalpina* Nagy, Upper Tithonian, Crassicollaria Zone, Massutiniana Subzone, **8** – sample 304-6, **9** – sample 307. **10-11** – *Calpionella elliptalpina* Nagy, Upper Tithonian, Crassicollaria Zone, Massutiniana Subzone, sample 310. **12-16** – *Calpionella alpina* Lorenz, Upper Tithonian, Crassicollaria Zone, Massutiniana Subzone, **12-13** – sample 307; Lower Berriasian, Calpionella Zone, Alpina Subzone, **14** – sample 313, **15** – sample 318, **16** – sample 321. **17-20** – *Calpionella minuta* Houša, Lower Berriasian, Calpionella Zone, Alpina Subzone, **17-18** – sample 318, **19-20** – sample 321. **21-22** – *Tintinnopsella doliphormis* (Colom), Lower Berriasian, Calpionella Zone, Alpina Subzone, **21** – sample 321, **22** – sample 322. **23-25** – *Remaniella ferasini* (Catalano), Lower Berriasian, Calpionella Zone, Remaniella Subzone, **23-24** – sample 323, **25** – sample 324. **26-28** – *Remaniella duranddelgai* Pop, Lower Berriasian, Calpionella Zone, Remaniella Subzone, **26-27** – sample 324, **28** – sample 325. **29-32** – *Remaniella colomi* Pop, Lower Berriasian, Calpionella Zone, Remaniella Subzone, **29** – sample 323, **31** – sample 325; Elliptica Subzone, **30, 32** – sample 326. **33** – *Remaniella filipescui* Pop, Lower Berriasian, Calpionella Zone, Elliptica Subzone, sample 327. **34** – *Tintinnopsella carpathica* (Murgeanu & Filipescu), Lower Berriasian, Calpionella Zone, Elliptica Subzone, sample 327. **35** – *Tintinnopsella longa* (Colom), Lower Berriasian, Calpionella Zone, Elliptica Subzone, sample 327. **36-37** – *Calpionella elliptica* Cadisch, Lower Berriasian, Calpionella Zone, Elliptica Subzone, **36** – sample 327, **37** – sample 328.




PLATE 3

Calpionellids of the Barlya section, Western Balkan Mts.

1-2 – *Calpionella elliptica* Cadisch, Lower Berriasian, Calpionella Zone, Elliptica Subzone, **1** – sample 328, **2** – sample 329. **3** – *Calpionella* sp.A, Lower Berriasian, Calpionella Zone, Elliptica Subzone, sample 328. **4** – *Calpionella minuta* Houša, Lower Berriasian, Calpionella Zone, Elliptica Subzone, sample 328. **5-8** – *Lorenziella hungarica* Knauer and Nagy, Lower Berriasian, Calpionella Zone, Elliptica Subzone, **5** – sample 330; Upper Berriasian, Calpionellopsis Zone, Simplex Subzone, **6** – sample 331, **7** – sample 332; Oblonga Subzone, **8** – sample 333. **9-11** – *Lorenziella plicata* Remane, Upper Berriasian, Calpionellopsis Zone, Oblonga Subzone, **9** – sample 333, **10-11** – sample 334. **12** – *Remaniella duranddelgai* Pop, Lower Berriasian, Calpionella Zone, Elliptica Subzone, sample 328. **13-16** – *Remaniella colomi* Pop, Lower Berriasian, Calpionella Zone, Elliptica Subzone, **13-14** – sample 328, **15-16** – sample 329-3. **17** – *Remaniella filipescui* Pop, Upper Berriasian, Calpionellopsis Zone, Oblonga Subzone, sample 333. **18** – *Remaniella cadischiana* (Colom), Lower Berriasian, Calpionella Zone, Elliptica Subzone, sample 329-3. **19-26** – *Calpionellopsis simplex* (Colom), Upper Berriasian, Calpionellopsis Zone, Simplex Subzone, **19** – sample 331, **20-22** – sample 332; Oblonga Subzone, **23-24** – sample 333, **25-26** – sample 334. **27-33** – *Calpionellopsis oblonga* (Cadisch), Upper Berriasian, Calpionellopsis Zone, Oblonga Subzone, **27-29** – sample 333, **30** – sample 334, **31-33** – sample 335. **34-35** – *Calpionellopsis* sp.A, Upper Berriasian, Calpionellopsis Zone, Oblonga Subzone, **34** – sample 333, **35** – sample 334. **36-37** – *Tintinnopsella subacuta* (Colom), Lower Berriasian, Calpionella Zone, Elliptica Subzone, **36** – sample 329-3, **37** – sample 330. **38-40** – *Tintinnopsella longa* (Colom), Upper Berriasian, Calpionellopsis Zone, Oblonga Subzone, **38** – sample 333, **39-40** – sample 334. **41-44** – *Tintinnopsella carpathica* (Murgeanu and Filipescu), Lower Berriasian, Calpionella Zone, Elliptica Subzone, **41** – sample 328, **42** – sample 329; Upper Berriasian, Calpionellopsis Zone, Simplex Subzone, **43** – sample 332; Oblonga Subzone, **44** – sample 334.




PLATE 4

Calpionellids of the Barlya section, Western Balkan Mts.

1-3 – *Calpionellopsis oblonga* (Cadisch), Upper Berriasian, Calpionellopsis Zone, Murgeanui Subzone, sample 336. **4-7** – *Calpionellopsis* sp.A, Upper Berriasian, Calpionellopsis Zone, Oblonga Subzone, **4-5** – sample 335; Murgeanui Subzone, **6-7** – sample 336. **8-9** – *Lorenziella plicata* Remane, Upper Berriasian, Calpionellopsis Zone, Murgeanui Subzone, **8** – sample 336; Lower Valanginian, Calpionellites Zone, Major Subzone, **9** – sample 341. **10** – *Remaniella filipes* Pop, Upper Berriasian, Calpionellopsis Zone, Murgeanui Subzone, sample 336. **11** – *Tintinnopsella subacuta* (Colom), Lower Valanginian, Calpionellites Zone, Darderi Subzone, sample 339. **12-15** – *Praecalpionellites murgeanui* (Pop), Upper Berriasian, Calpionellopsis Zone, Murgeanui Subzone, **12** – sample 336, **13-14** – sample 337; Lower Valanginian, Calpionellites Zone, Darderi Subzone, **15** – sample 340. **16-17** – *Praecalpionellites siriniaensis* Pop, Lower Valanginian, Calpionellites Zone, Darderi Subzone, **16** – sample 338; Major Subzone, **17** – sample 343. **18-19** – *Praecalpionellites hillebrandti* Grün & Blau, Lower Valanginian, Calpionellites Zone, Darderi Subzone, **18** – sample 339, **19** – sample 338. **20-24** – *Calpionellites darderi* (Colom), Lower Valanginian, Calpionellites Zone, Darderi Subzone, **20-21** – sample 338, **22-23** – sample 340; Major Subzone, **24** – sample 343. **25-36** – *Calpionellites major* (Colom), Lower Valanginian, Calpionellites Zone, Major Subzone, **25** – sample 341, **26-27** – sample 342, **28-30** – sample 343, **31-36** – sample 344. **37-38** – *Calpionellites coronatus* Trejo, Lower Valanginian, Calpionellites Zone, Darderi Subzone, sample 338. **39-44** – *Calpionellites caravacaensis* Allemann, Lower Valanginian, Calpionellites Zone, Major Subzone, **40** – sample 342, **41-42** – sample 343, **39, 43-44** – sample 344. **45-46** – *Sturiella* aff. *oblonga* Borza, Upper Berriasian, Calpionellopsis Zone, Murgeanui Subzone, **45** – sample 337; Lower Valanginian, Calpionellites Zone, Major Subzone, **46** – sample 344. **47-49** – *Tintinnopsella carpathica* (Murgeanu & Filipescu), Upper Berriasian, Calpionellopsis Zone, Murgeanui Subzone, **47-48** – sample 336; Lower Valanginian, Calpionellites Zone, Darderi Subzone, **49** – sample 340. **50-53** – *Tintinnopsella longa* (Colom), Lower Valanginian, Calpionellites Zone, Darderi Subzone, **50-51** – sample 338, **52** – sample 339; Major Subzone, **53** – sample 341. **54** – *Tintinnopsella dacica* Filipescu & Dragastan, Lower Valanginian, Calpionellites Zone, Major Subzone, sample 341.




PLATE 5

Calpionellids of the Gintsi 1 and Gintsi 2 sections, Western Balkan Mts.

1-5 – *Daciella svinitensis* Pop, Gintsi 1 section, Lower Tithonian, Chitinoidea Zone, Dobeni Subzone, sample Gin-011. **6-7** – *Borziella slovenica* (Borza), Gintsi 1 section, Lower Tithonian, Chitinoidea Zone, Dobeni Subzone, **6** – sample Gin-011; Gintsi 2 section, **7** – sample 610. **8** – *Carpathella rumanica* Pop, Gintsi 2 section, Lower Tithonian, Chitinoidea Zone, Dobeni Subzone, sample 610. **9-13** – *Daciella danubica* Pop, Gintsi 1 section, Lower Tithonian, Chitinoidea Zone, Dobeni Subzone, sample Gin-011. **14-15** – *Dobeniella colomi* (Borza), Gintsi 1 section, Lower Tithonian, Chitinoidea Zone, Dobeni Subzone, sample Gin-011. **16** – *Daciella* sp. A, Gintsi 1 section, Lower Tithonian, Chitinoidea Zone, Dobeni Subzone, sample Gin-011. **17-20** – *Longicollaria dobeni* (Borza), Gintsi 1 section, Lower Tithonian, Chitinoidea Zone, Dobeni Subzone, **17** – sample Gin-08, **18-19** – sample Gin-011, **20** – sample Gin-09. **21-23** – *Chitinoidea boneti* Doben, Gintsi 1 section, Upper Tithonian, Chitinoidea Zone, Boneti Subzone, **21** – sample Gin-03, **22** – sample Gin-02, **23** – sample Gin-01. **24-25** – *Chitinoidea elongata* Pop, Gintsi 2 section, Upper Tithonian, Chitinoidea Zone, Boneti Subzone, **24** – sample 611; Gintsi 1 section, **25** – sample Gin-02. **26** – *Longicollaria insueta* (Řehánek), Gintsi 1 section, Upper Tithonian, Chitinoidea Zone, Boneti Subzone, sample Gin-01. **27-28** – *Dobeniella bermudezi* (Furrazola-Bermúdez), Gintsi 1 section, Upper Tithonian, Chitinoidea Zone, Boneti Subzone, **27** – sample Gin-03, **28** – sample Gin-02. **29-32** – *Dobeniella cubensis* (Furrazola-Bermúdez), Gintsi 2 section, Upper Tithonian, Chitinoidea Zone, Boneti Subzone, **29** – sample 611; Gintsi 1 section, **30** – sample Gin-02, **31-32** – sample Gin-1. **33-35** – *Praetintinnopsella andrusovi* Borza, Gintsi 1 section, Upper Tithonian, Praetintinnopsella Zone, sample Gin-3. **36-37** – *Tintinnopsella* aff. *carpathica* (Murgeanu & Filipescu), Gintsi 1 section, Upper Tithonian, Praetintinnopsella Zone, sample Gin-4. **38-41** – *Tintinnopsella carpathica* (Murgeanu & Filipescu), Gintsi 1 section, Upper Tithonian, Praetintinnopsella Zone, **38** – sample Gin-3; Crassicollaria Zone, Remanei Subzone, **39-40** – sample Gin-7, **41** – sample Gin-5. **42-43** – *Tintinnopsella remanei* Borza, Gintsi 1 section, Upper Tithonian, Crassicollaria Zone, Remanei Subzone, sample Gin-5. **44-46** – *Crassicollaria intermedia* (Durand-Delga), Gintsi 1 section, Upper Tithonian, Crassicollaria Zone, Remanei Subzone, **44-45** – sample Gin-7, **46** – sample Gin-9. **47-48** – *Crassicollaria massutiniana* (Colom), Gintsi 1 section, Upper Tithonian, Crassicollaria Zone, Massutiniana Subzone, **47** – sample Gin-10, **48** – sample Gin-11. **49** – *Crassicollaria parvula* Remane, Gintsi 1 section, Upper Tithonian, Crassicollaria Zone, Massutiniana Subzone, sample Gin-10. **50-51, 53-54** – *Calpionella grandalpina* Nagy, Gintsi 1 section, Upper Tithonian, Crassicollaria Zone, Massutiniana Subzone, **50-51** – sample Gin-10, **53-54** – sample Gin-11. **52** – *Calpionella alpina* Lorenz, Gintsi 1 section, Upper Tithonian, Crassicollaria Zone, Massutiniana Subzone, sample Gin-10.




PLATE 6

Calpionellids of the Gintsi 1 and Gintsi 2 sections, Western Balkan Mts.

1 – *Calpionella grandalpina* Nagy, Gintsi 1 section, Upper Tithonian, Crassicollaria Zone, Massutiniana Subzone, sample Gin-11. **2** – *Calpionella elliptalpina* Nagy, Gintsi 1 section, Upper Tithonian, Crassicollaria Zone, Massutiniana Subzone, sample Gin-14. **3-7** – *Calpionella alpina* Lorenz, Gintsi 1 section, Upper Tithonian, Crassicollaria Zone, Massutiniana Subzone, **3** – sample Gin-11; Lower Berriasian, Calpionella Zone, Alpina Subzone, **4-6** – sample Gin-19; Remaniella Subzone, **7** – sample Gin-30. **8-9** – *Calpionella minuta* Houša, Gintsi 1 section, Lower Berriasian, Calpionella Zone, Remaniella Subzone, **8** – sample Gin-30, **9** – sample Gin-31. **10-13** – *Calpionella* sp.A, Gintsi 1 section, Lower Berriasian, Calpionella Zone, Remaniella Subzone, **10-11** – sample Gin-30, **12** – sample Gin-31; Elliptica Subzone, **13** – sample Gin-35. **14** – *Calpionella elliptica* Cadisch, Gintsi 2 section, Lower Berriasian, Calpionella Zone, Elliptica Subzone, sample 11. **15-17** – *Crassicollaria massutiniana* (Colom), Gintsi 1 section, Upper Tithonian, Crassicollaria Zone, Massutiniana Subzone, sample Gin-11. **18-19** – *Crassicollaria brevis* Remane, Gintsi 1 section, Upper Tithonian, Crassicollaria Zone, Massutiniana Subzone, sample Gin-14. **20-29** – *Crassicollaria parvula* Remane, Gintsi 1 section, Upper Tithonian, Crassicollaria Zone, Massutiniana Subzone, **20-22** – sample Gin-11, **23-24** – sample Gin-12; Lower Berriasian, Calpionella Zone, Alpina Subzone, **25-28** – sample Gin-19; Remaniella Subzone, **29** – sample Gin-30. **30-31** – *Crassicollaria colomi* Doben, Gintsi 1 section, Lower Berriasian, Calpionella Zone, Alpina Subzone, sample Gin-19. **32** – *Tintinnopsella carpathica* (Murgeanu & Filipescu), Gintsi 1 section, Lower Berriasian, Calpionella Zone, Alpina Subzone, sample Gin-19. **33-35** – *Tintinnopsella doliphormis* (Colom), Gintsi 1 section, Lower Berriasian, Calpionella Zone, Remaniella Subzone, **33** – sample Gin-30; Gintsi 2 section, Lower Berriasian, Calpionella Zone, Alpina Subzone, **34** – sample 8b; Elliptica Subzone, **35** – sample 12. **36-41** – *Remaniella ferasini* (Catalano), Gintsi 1 section, Lower Berriasian, Calpionella Zone, Remaniella Subzone, **37** – sample Gin-31, **38** – sample Gin-32, **39-41** – sample Gin-33; Gintsi 2 section, Lower Berriasian, Calpionella Zone, Remaniella Subzone, **36** – sample 10. **42-49** – *Remaniella duranddelgai* Pop, Gintsi 1 section, Lower Berriasian, Calpionella Zone, Remaniella Subzone, **42-44** – sample Gin-30, **45** – sample Gin-31, **46-47** – sample Gin-32, **48-49** – sample Gin-33. **50-51** – *Remaniella colomi* Pop, Gintsi 2 section, Lower Berriasian, Calpionella Zone, Elliptica Subzone, **50** – sample 11, **51** – sample 18. **52-55** – *Remaniella catalanoi* Pop, Gintsi 1 section, Lower Berriasian, Calpionella Zone, Remaniella Subzone, **52** – sample Gin-33; Elliptica Subzone, **53-55** – sample Gin-34.




PLATE 7

Calpionellids of the Gintsi 1 and Gintsi 2 sections, Western Balkan Mts.

1-2 – *Calpionellopsis simplex* (Colom), Gintsi 1 section, Upper Berriasian, Calpionellopsis Zone, Murgeanui Subzone, **1** – sample Gin-36; Gintsi 2 section, Upper Berriasian, Calpionellopsis Zone, Murgeanui Subzone, **2** – sample 21. **3-6** – *Calpionellopsis oblonga* (Cadisch), Gintsi 2 section, Upper Berriasian, Calpionellopsis Zone, Murgeanui Subzone, **3-4** – sample 19, **5-6** – sample 20. **7-8** – *Calpionellopsis* sp.A, Gintsi 2 section, Upper Berriasian, Calpionellopsis Zone, Murgeanui Subzone, **7** – sample 19, **8** – sample 21. **9-12** – *Lorenziella hungarica* Knauer & Nagy, Gintsi 2 section, Lower Valanginian, Calpionellites Zone, Major Subzone, **9** – sample 28, **10** – sample 30, **11** – sample 35, **12** – sample 38. **13-14** – *Lorenziella plicata* Remane, Gintsi 2 section, Upper Berriasian, Calpionellopsis Zone, Murgeanui Subzone, **13** – sample 20; Lower Valanginian, Calpionellites Zone, Major Subzone, **14** – sample 35. **15-17** – *Praecalpionellites murgeanui* (Pop), Gintsi 2 section, Upper Berriasian, Calpionellopsis Zone, Murgeanui Subzone, **15** – sample 21; Lower Valanginian, Calpionellites Zone, Darderi Subzone, **16-17** – sample 27. **18-20** – *Praecalpionellites siriniaensis* Pop, Gintsi 2 section, Upper Berriasian, Calpionellopsis Zone, Murgeanui Subzone, **18** – sample 19; Lower Valanginian, Calpionellites Zone, Darderi Subzone, **19** – sample 26; Major Subzone, **20** – sample 36. **21** – *Praecalpionellites* sp. A, Gintsi 2 section, Lower Valanginian, Calpionellites Zone, Darderi Subzone, sample 26. **22** – *Remaniella borzai* Pop, Gintsi 2 section, Lower Valanginian, Calpionellites Zone, Major Subzone, sample 34. **23-24** – *Remaniella cadischiana* (Colom), Gintsi 2 section, Upper Berriasian, Calpionellopsis Zone, Murgeanui Subzone, **23** – sample 19, **24** – sample 20. **25-26** – *Calpionellites darderi* (Colom), Gintsi 2 section, Lower Valanginian, Calpionellites Zone, Darderi Subzone, **25** – sample 27; Major Subzone, **26** – sample 32. **27-29** – *Calpionellites coronatus* Trejo, Gintsi 2 section, Lower Valanginian, Calpionellites Zone, Major Subzone, sample 36. **30-32** – *Calpionellites major* (Colom), Gintsi 2 section, Lower Valanginian, Calpionellites Zone, Major Subzone, **30** – sample 28, **31** – sample 29, **32** – sample 30. **33-35** – *Calpionellites caravacaensis* Allemann, Gintsi 2 section, Lower Valanginian, Calpionellites Zone, Major Subzone, **33-34** – sample 34, **35** – sample 37. **36-41** – *Tintinnopsella carpathica* (Murgeanu & Filipescu), Gintsi 2 section, Upper Berriasian, Calpionellopsis Zone, Murgeanui Subzone, **36-37** – sample 20; Lower Valanginian, Calpionellites Zone, Major Subzone, **38** – sample 33, **39-40** – sample 36, **41** – sample 37. **42-44** – *Tintinnopsella subacuta* (Colom), Gintsi 2 section, Lower Valanginian, Calpionellites Zone, Darderi Subzone, **42** – sample 27; Major Subzone, **43** – sample 35, **44** – sample 36. **45-48** – *Tintinnopsella longa* (Colom), Gintsi 2 section, Upper Berriasian, Calpionellopsis Zone, Murgeanui Subzone, **45** – sample 19; Lower Valanginian, Calpionellites Zone, Major Subzone, **46** – sample 35, **47** – sample 38, **48** – sample 39. **49-50** – *Sturiella oblonga* Borza, Gintsi 2 section, Lower Valanginian, Calpionellites Zone, Major Subzone, **49** – sample 28, **50** – sample 33. **51-53** – *Tintinnopsella dacica* Filipescu & Dragastan, Gintsi 2 section, Lower Valanginian, Calpionellites Zone, Major Subzone, **51-52** – sample 29, **53** – sample 33.

